

RESEARCH ARTICLE

Open Access



# Metabolomic changes in polyunsaturated fatty acids and eicosanoids as diagnostic biomarkers in *Mycobacterium avium* ssp. *paratuberculosis* (MAP)-inoculated Holstein–Friesian heifers

Emma N. Taylor<sup>1</sup>, Manfred Beckmann<sup>1</sup>, Glyn Hewinson<sup>1,2</sup>, David Rooke<sup>3</sup>, Luis A. J. Mur<sup>1\*</sup> and Ad P. Koets<sup>4,5\*</sup> 

## Abstract

*Mycobacterium avium* subspecies *paratuberculosis* (MAP) is the causative organism of Johne's disease, a chronic granulomatous enteritis of ruminants. We have previously used naturally MAP-infected heifer calves to document metabolomic changes occurring in MAP infections. Herein, we used experimentally MAP-inoculated heifer calves to identify biomarkers for MAP infections. At 2-weeks of age, 20 Holstein–Friesian (HF) calves were experimentally inoculated with MAP. These calves, along with 20 control calves, were sampled biweekly up to 13-months of age and then monthly up to 19-months of age. Sera were assessed using flow infusion electrospray high-resolution mass spectrometry (FIE-HRMS) on a Q Exactive hybrid quadrupole-Orbitrap mass spectrometer for high throughput, sensitive, non-targeted metabolite fingerprinting. Partial least squares-discriminate analysis (PLS-DA) and hierarchical cluster analysis (HCA) discriminated between MAP-inoculated and control heifer calves. Out of 34 identified metabolites, six fatty acyls were able to differentiate between experimental groups throughout the study, including 8, 11, 14-eicosatrienoic acid and cis-8, 11, 14, 17-eicosatetraenoic acid which were also detected in our previous study and so further suggested their value as biomarkers for MAP infection. Pathway analysis highlighted the role of the alpha-linoleic acid and linoleic acid metabolism. Within these pathways, two broad types of response, with a rapid increase in some saturated fatty acids and some n-3 polyunsaturated fatty acids (PUFAs) and later n-6 PUFAs, became predominant. This could indicate an initial anti-inflammatory colonisation phase, followed by an inflammatory phase. This study demonstrates the validity of the metabolomic approach in studying MAP infections. Nevertheless, further work is required to define further key events, particularly at a cell-specific level.

**Keywords:** *Mycobacterium avium* subsp. *paratuberculosis*, metabolomics, alpha-linoleic acid and linoleic acid metabolism, polyunsaturated fatty acids, eicosanoids

## Introduction

Johne's disease, caused by *Mycobacterium avium* subspecies *paratuberculosis* (MAP), is a chronic granulomatous enteritis of ruminants. MAP is frequently transmitted via the ingestion of colostrum, milk and faeces [1]. Following exposure to MAP, infected cattle enter a prolonged incubation period prior to the onset of shedding and clinical

\*Correspondence: lum@aber.ac.uk; ad.koets@wur.nl

<sup>1</sup> Institute of Biological, Environmental and Rural Sciences (IBERS), Aberystwyth University, Ceredigion SY23 3DA, UK

<sup>4</sup> Wageningen Bioveterinary Research, 8221 RA Lelystad, The Netherlands  
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

signs, such as; weight loss, diarrhoea and reduced milk yields [2]. With estimated herd prevalence's exceeding 50% in countries with intensive dairy production systems [3] and 10-year annual losses of ~\$49 million (~£35 million) in the UK [4], MAP represents an on-going economic and animal welfare challenge for farmers.

Identifying MAP-infected young stock is particularly challenging due to an absence of clinical signs [5], and detection being based on bacterial shedding patterns that are influenced by the level and age of calves at MAP exposure [6]. Individual faecal PCR tests and serum ELISAs are commonly used to identify MAP-infected cattle [7] but exhibit poor sensitivity values of approximately 4% and 13% in light-to-moderate faecal shedders [8]. Furthermore, interferon- $\gamma$  tests only detect approximately half of MAP faecal culture positive cattle aged between 12 and 23 months of age [9] and milk ELISA tests are limited to post-partum use only. Consequently, a diagnostic tool capable of detecting MAP-infected youngstock would benefit farmers worldwide.

Omic approaches encapsulate various subdivisions of systems biology, including genomics, proteomics, lipidomics and metabolomics [10]. The application of these approaches in MAP research has highlighted multiple potential biomarkers, but these findings require further validation [11–13]. Proteomic approaches identified antigenically distinct recombinant MAP cell envelope proteins; Sdh A and MAP1233m and their detection using ELISA showed maximum sensitivity and specificity values of 94% and 100%, respectively [12]. Further lipidomic approaches were used to show that glycopeptidolipids are replaced by lipopentapeptides within the MAP cell wall. Subsequently, a lipopentapeptide L5P-based test exhibited a sensitivity of 82% [11]. These studies suggest that the performance of these omic study-derived biomarkers exceeds many commercially available tests [8]. This stated, these have focused on post-partum cattle [12, 13] so that the application of these biomarkers in youngstock is questionable.

In contrast, metabolomic studies have focused on both calves and young stock. Analysis of sera from experimentally MAP-infected calves using  $^1\text{H}$  nuclear magnetic resonance (NMR) spectrometry showed metabolomic changes indicative of energy shortages, elevated lipid metabolism and increased protein turnover [14]. Similar energy and lipid-related changes were reported within the sera of mature, naturally MAP-infected cows following analysis via direct analysis in real time coupled with high resolution mass spectrometry (DART-HRMS) [15]. However, the metabolites identified by de Buck et al. [14] exhibited similar levels between MAP-infected and control cattle until 200 days post-infection. Likewise, the metabolites identified by Tata et al. [15] showed

overlapping levels between infected, infectious and control groups. Thus, the targeted metabolites were unsuitable as biomarkers for MAP infections.

We have previously examined the sera metabolite profiles of naturally MAP-infected Holstein–Friesian (HF) heifer calves compared to controls between 1 and 19 months of age and identified 33 metabolites which were differentially accumulated in MAP-infected heifer calves [16]. Crucially, five of these metabolites were elevated throughout the study. However, to more clearly examine the time dependent changes following MAP infection in youngstock, we herein use metabolomics to assess a complementary experiment based on HF heifer calves experimentally inoculated with MAP. Biweekly sera samples of heifer calves up to 13 months of age and monthly sera samples of heifer calves up to 19 months of age were analysed using flow infusion electrospray high-resolution mass spectrometry (FIE-HRMS).

## Materials and methods

### Animal samples

The animal experiments were approved by the Animal Welfare Body of Utrecht University (permit number 0202.0806) in accordance with the Dutch regulations on animal experimentation. All samples were bio-archived serum samples previously described in Koets et al. [17].

Sera were obtained from 20 experimentally MAP-inoculated, and 20 controls heifer calves, sampled biweekly until 12 months of age and by monthly until 19 months of age. Control HF heifer calves were sourced at approximately 2 weeks of age from MAP-negative herds. The MAP status of control farms was confirmed via faecal culture from cattle  $\geq 24$  months of age every 6 months or 12 months for a minimum of 8 years as part of a Dutch national MAP surveillance program. All MAP-inoculated heifer calves were confirmed to be MAP positive in at least one tissue sample at necropsy. No macroscopic lesions were observed.

MAP-inoculated calves were orally inoculated with approximately  $1.8 \times 10^4$  colony forming units (cfu), supplied across nine doses of 20 g of faeces, and mixed with 100 mL milk replacer over 21 days [17]. The faeces were derived from a MAP-infected cow, whose infection status was confirmed via faecal culture and PCR. Faecal samples were collected from all heifer calves 0-, 14-, 126-, 280-, 406- and 532- days post-inoculation. Of the MAP-inoculated heifer calves, 13 were MAP faecal culture positive and 7 were negative, all control heifer calves were culture negative. MAP faecal culture positive heifer calves displayed faecal shedding at either 14 (one heifer calf) or 126 (twelve heifer calves) days post-inoculation. Faecal samples from heifer calves were processed using the routine diagnostic MAP faecal culture system at the

Veterinary Health Service laboratory (Royal GD), Deventer, the Netherlands.

Heifer calves were screened using a commercially available absorbed ELISA assay following the manufacturer's instructions (Institute Pourquier, Montpellier, France) using blood samples collected 0-, 14-, 126-, 280-, 406- and 532- days post-inoculation. Samples with a sample to negative (S/N) ratio > 59% were considered MAP positive. Of the MAP-inoculated calves, 8 were ELISA positive and 12 were negative. Of the control calves, 2 were ELISA positive and 18 were negative. All positive ELISA results occurred a minimum of 406 days post-inoculation.

#### Untargeted metabolite fingerprinting by flow infusion electrospray ionization high resolution mass spectrometry (FIE-HRMS)

Sera were prepared as described by Beckmann et al. [18] with minor amendments. Samples were defrosted on ice, vortexed for 5 s and 200  $\mu$ L was pipetted into 1520  $\mu$ L pre-chilled solvent mix (methanol/chloroform [4/1 v/v]) containing 1 micro-spoon of glass beads glass beads (<106  $\mu$ m diameter, Sigma, UK). Samples were then vortexed for 5 s, shaken for 15 min at +4  $^{\circ}$ C and kept at -20  $^{\circ}$ C for 20 min. Following centrifugation at 21 000 $\times$ g and 4  $^{\circ}$ C for 5 min, 100  $\mu$ L of the plasma supernatant was transferred into mass spectrometry vials along with 100  $\mu$ L methanol/water [70/30 v/v]. Samples were stored at -80  $^{\circ}$ C until analysis using flow infusion electrospray ionization high-resolution mass spectrometry (FIE-HRMS). For each sample, 20  $\mu$ L were injected into a flow of 60  $\mu$ L per minute water-methanol, at a ratio of 70% water and 30% methanol, using a Surveyor flow system into a Q Exactive plus mass analyser instrument with UHPLC system (Thermo Fisher Scientific $^{\circ}$ , Bremen, Germany) for high throughput FIE-HRMS. Data acquisition for each serum sample was done by alternating the positive and negative ionisation modes, throughout four different scan ranges (15–110 m/z, 100–220 m/z, 210–510 m/z, 500–1200 m/z) with an acquisition time of 2 min.

#### Statistical analysis

Metabolomic data were analysed using MetaboAnalyst 4 [19]. Data were subjected to interquartile range-based filtering,  $\log_{10}$  transformations and Pareto scaling. Time series analyses used false-discovery rate (FDR) adjusted two-way ANOVA tests to identify m/z which significantly ( $p$ -values < 0.05) differed between experimental classes. Partial least squares-discriminate analysis (PLS-DA) were used to visualise the differences between the experimental classes. Variables of importance for the projection (VIP) scores (> 1) following multivariate analyses were also used to indicate m/z which discriminated

between the classes. Random forest (RF) was used as an alternative multivariate classification test. The major sources of variation were displayed using unsupervised hierarchical clustering analysis (HCA). Area under the curve (AUC) assessments, based on sensitivity and specificity estimates, were used to determine the accuracy of the target m/z as potential biomarkers.



Significant m/z were identified based on accurate mass (5 ppm) using the DIMEdb database [20] based on their ionised masses, molecular formula and the Bovine Metabolome Database [21]. All isotopes/adducts were considered. Metabolite set enrichment analysis (MSEA) using over representation analysis (ORA) was used to highlight key biochemical pathways.

#### Results

Serum samples were profiled by FIE-HRMS and the derived spectra assessed by PLS-DA. PLS-DA of the data from both positive and negative ionisations indicated a clear differentiation between MAP-inoculated and control heifer calves (Figure 1A). The sources of variation most utilised in the PLS-DA were indicated using a VIP plot (>1.0) based on two components (Figure 1B). This indicated that the eicosanoid, 8,11,14 eicosatrienoic acid (C20:4) was the most notable discriminator between MAP-inoculated and control heifer calves. Indeed, other eicosanoids were also shown to have fairly high VIP scores, for example, eicosenoic acid (20:1) and 6Z, 9Z-octadecadienoic acid (C18:2).

Subsequently, the derived metabolomes were assessed using time series analysis, adjusted for FDR, two-way ANOVA. Significant metabolites were identified, and all were significantly affected by time and MAP inoculation status $\times$ time ( $p$  < 0.05) (Additional file 1). Out of 34 identified metabolites, 25 were fatty acyls and notable subclass included; fatty acids and conjugates (16), eicosanoids (6) and linoleic acid and derivatives (3) (Additional file 2).

Significantly altered metabolite changes were visualised using a heat map (Figure 2). The accumulation of identified metabolites was significantly higher in MAP-inoculated heifer calves compared to controls. However, the timings of metabolite accumulations differed between clades, whereby clades are defined as groups of metabolites which demonstrate comparable levels over time. Some changes were seen quickly following MAP inoculation (within 1 month) and remained elevated throughout the timeframe of the experiment. These metabolites are included in Figure 2 clade 3 and included alpha linolenic acid, 8,11,14 eicosatrienoic acid and stearic acid. The elevation of other metabolites (Figure 2, clade 4) was equally rapid following MAP inoculation but started to decrease after 12 months. Examples of metabolites in these categories included the fatty acids; myristic and palmitic acid,



as well as 6Z, 9Z-octadecadienoic acid (C18:2). Other metabolites were only elevated after 6 to 12 months and then remained higher (Figure 2, clade 1) and include

important eicosanoids, such as prostaglandin E1 and leukotriene B4. The eicosanoid, thromboxane A2, was more rapidly elevated and was found in Figure 2, clade 3.



Interestingly, many eicosanoid levels in the sera of control heifer calves shown in Figure 2 clade 1, exhibited some slight increases over time, possibly due to heifer growth and development.

To better visualise the biological events indicated by these metabolomic changes, identified metabolites were related to different pathways by MSEA using ORA analysis. This indicated that alpha-linoleic acid and linoleic acid metabolism was the only pathway to be significantly ( $p < 0.05$ ) affected by MAP inoculation status (Additional file 3). This pathway is depicted schematically in Figure 3, where the metabolites identified in our study are indicated. To further illustrate changes in the pathway, the accumulation patterns of individual metabolites (8, 11, 14-eicosatrienoic acid, alpha-linoleic acid, *cis*-8, 11, 14, 17-eicosatetraenoic acid and eicosapentaenoic acid) are plotted (Figure 4). Thus, elevated levels of alpha-linoleic acid derivatives was linked to increased levels of docosahexaenoic acid. This metabolite is shown in Figure 2, clade 4, indicating that its levels decreased >11 months following MAP infection. However, the linoleic acid derivative pathway, linked to cyclooxygenase (COX) and lipoxygenase (LOX)-dependent pathway, was active as shown by the detection of many prostaglandin, thromboxane and leukotriene derivatives. These metabolites increased more slowly following MAP inoculation (Figure 2, clade 1).

Excluding docosahexanoic acid, whereby the interquartile ranges of experimental groups overlapped 15 months post MAP inoculation (Additional file 4), metabolites identified within the alpha-linoleic acid and linolenic acid metabolism showed no evidence of increases in control heifer calves over time (Figure 4), suggesting that these changes were intimately associated with MAP inoculation. It was also noted that, the linoleic pathway metabolites, 10-octadecenoic acid and heptadecanoic acid showed no increases in control heifer calves over time (Additional file 5). Consequently, RF and AUC assessments were used to determine the potential of these six metabolites as biomarkers for MAP infections. RF assessments across all ages indicated classification errors for MAP-inoculated and control heifer calves of only 0.00781 and 0.00313, respectively (Table 1). Moreover, AUC assessments of these metabolites at 19-months of age suggested individual metabolite AUC values of 1.0 (C.I 0.913–0.999) (Additional file 2) and a combined AUC value of 0.999 (C.I 0.992–1.000), as well as sensitivity and specificity values of 99.2% and 99.6%, irrespective of age.

Additionally, we compared the ability of identified metabolites, faecal culture and serum ELISA tests to differentiate between experimental groups. Although our metabolites were unable to differentiate between heifer


calves based on faecal culture or serum ELISA status (Figures 5 and 6), our metabolites better discriminated between experimental groups based on MAP inoculation status.

## Discussion

Johne's disease, caused by MAP, has a significant impact on cattle health alongside severe economic consequences. Current diagnostic tests for MAP, including faecal PCR, serum ELISA [8] and IFN- $\gamma$  tests [9] rely on detecting a cell mediated immune (CMI) response, antibody production or bacterial shedding. As a result, they perform poorly between the initial CMI response and the sub (clinical) stages of infection [22]. The application of omic approaches have suggested that they could be a potential source of new biomarkers with sensitivity and specificity values reported ranging between 82 and 94% [11, 12]. However, no omic study has assessed MAP infection associated changes in youngstock. Our previous metabolomic study identified 33 metabolites which were differentially accumulated within naturally MAP-infected heifer calves, including 5 metabolites which were elevated throughout the study [16]. In this study, we examined experimentally MAP-inoculated heifer calves to more clearly assess early changes occurring following infection and to validate the previously observed metabolites changes in an analogous study.

Our findings identified 11 metabolites in MAP-inoculated heifer calves that were also observed in our Taylor et al. [16] study which focused on natural MAP infections. The prominence of metabolites from the fatty acyl class and fatty acids and conjugates subclass, alongside the eicosanoid subclass was notable in both studies (Additional file 2) [16]. However, there was an absence of metabolites belonging to the amino acids, peptides and analogues subclass in our current study (Additional file 2). Changes to bioenergetic and uraemic pathways also seem specific to naturally infected heifer calves [16]. These changes may be attributable to differences in bacterial load, as indicated by the faecal culture results. Taylor et al. [16] noted MAP-positive faecal culture results within all MAP-positive heifers. In contrast, only 13 out of 20 MAP-inoculated heifer calves were MAP faecal culture positive within our current study. Thus, implying that a higher bacterial load is required to observe bioenergetic and uraemic changes, compared to fatty acyl related changes. In contrast, fatty acyls changes appear to be consistent in both naturally and experimentally MAP inoculated heifer calves studies. This could support their importance as potential immunomodulators.


Moreover, although Taylor et al. [16] reported that accumulations of leukotriene B<sub>4</sub>, bicyclo prostaglandin E<sub>2</sub>, itaconic acid, 2-hydroxyglutaric acid and



**Figure 3** The metabolomic impact of MAP on selected metabolites. Metabolites include; selected unsaturated fatty acids, metabolites located within alpha-linolenic acid and linoleic acid metabolism and selected eicosanoids. Coloured boxes = identified metabolites, white boxes = metabolites which were not identified. *FADS1* = fatty acid desaturase 1, *FADS2* = fatty acid desaturase 2, *ELOV5* = elongation of very long chain fatty acids protein 5, *COX* = cyclooxygenase, *5-LOX* = 5-lipoxygenase, PGH<sub>2</sub> = prostaglandin H<sub>2</sub>, 5-HPETE = 5-hydroperoxyeicosatetraenoic acid, 5-HETE = 5-hydroxyeicosatetraenoic acid, TXA<sub>2</sub> = carbocyclic thromboxane A<sub>2</sub> LTA<sub>4</sub> = leukotriene A<sub>4</sub>, LTB<sub>4</sub> = leukotriene B<sub>4</sub>, 10, 11-LTB<sub>4</sub> = 10,11-dihydro-leukotriene B<sub>4</sub>.

N6-acetyl-L-lysine were significantly elevated throughout the study, accumulations of the majority of metabolites fluctuated within MAP infected and control heifers up to

13 months of age. In our current study, eicosanoids also followed this pattern (clade 1, Figure 2). However, fatty acids and conjugates demonstrated minimal fluctuations




**Table 1** Cross validation results of the random forest assessments. These assessments classified serum samples from heifers between 1-month and 19-months of age as MAP-inoculated or control based on metabolite levels of 8,11,14-eicosatrienoic acid, 10-octadecenoic acid, alpha-linolenic acid, cis-8,11,14,17-eicosatetraenoic acid, eicosapentaenoic acid and heptadecanoic acid

|                | MAP-inoculated | Control | Class error |
|----------------|----------------|---------|-------------|
| MAP-inoculated | 638            | 2       | 0.00313     |
| Control        | 5              | 635     | 0.00781     |

(clade 3, Figure 2). These observations may reflect MAP inoculated heifer calves receiving nine doses of 20 g of faeces, as opposed to being naturally infected. Fecteau et al. [23] and Kaevska et al. [24] observed significant differences within the faecal microbiota of MAP shedding

cattle, compared to non-shedding cattle. Likewise, Derakhshani et al. [25] noted the influence of rumen microbiota on functional pathways, such as; carbohydrate, lipid and amino acid metabolism. More recently, Umanets et al. [26] demonstrated how intestinal microbiota can be utilised to predict the severity of future MAP shedding with AUC accuracies between 0.91 and 0.92 for cattle aged between 12 and 24 months of age. Thus, repeat inoculations of MAP culture positive faeces may aid the establishment of a MAP permissive microbiome.


Our previous examination of MAP effects on the sera in naturally infected heifer calves, had indicated the prominence of fatty acid changes [16] and this was again the case in this study. Accumulations of palmitic acid (C16:0) and stearic acid (C18:0) were observed in naturally and experimentally MAP-inoculated heifer calves (Figure 2). Such an observation was partially in agreement with those of Tata et al. [15] who also



targeted palmitic acid as a MAP-infection associated change although not stearic acid. In another MAP focused metabolomic study, neither of these metabolites were identified [14]. Although, this could be seen to reduce the validity of our observations, these differences may be attributable to a mixture of biological factors, such as; MAP dose and bacterial strain, and our use of a direct infusion MS approach, as opposed to NMR spectrometry. Equally, there is independent corroboration of our observations in a transcriptional assessment of MAP-infected sheep. Thus, the expression of FABP3 and PDK4, which are associated with the

oxidation of palmitic acid, as well as, FABP3 and ORL1, which are linked to the uptake of palmitic acid, were elevated in paucibacillary and resilient MAP-infected sheep, respectively [27]. It was not possible to assess how far these changes reflected metabolic events occurring in either the heifer calves or in the MAP organisms or both combined. It should be noted that palmitic acid and stearic acid are common biochemical features of *Mycobacterium* [28, 29]. To define the relative origins of particular metabolites will require the use of such as isotopically labelled MAP in further infection experiments as demonstrated in other pathosystems [30].





**Figure 6** HCA of the metabolite changes alongside serum ELISA results at 15-months of age. The metabolite changes are those which differentiate between MAP-inoculated and control heifer calves in the combined ionisation mode *m/z*.

The likely increased depth of the direct infusion MS approach could have led our targeting of the alpha-linolenic acid (C18:3) and linoleic acid (C18:2) metabolism. This pathway was not identified in other MAP-focused metabolomic studies [14, 15] but was significantly affected by MAP inoculation status in our current (Additional file 3) and previous study [16]. The metabolites that formed part of the alpha-linolenic acid/linoleic acid (C18:2) pathway were previously suggested to be potential biomarkers for MAP infection in heifer calves [16]. This underlined the validity of changes in 8, 11, 14-eicosatrienoic acid, cis-8, 11, 14–17-eicosatetraenoic acid,

eicosapentaenoic acid and docosahexaenoic acid as early biomarkers for MAP infection. Two patterns of alpha-linolenic acid/linoleic acid metabolite accumulation was observed. Most metabolites in this pathway increased over time (Figure 2), however, docosahexaenoic acid showed a rapid increase, followed by a decline in levels after ~10 months. This raises the question as to the clinical significance of these differential accumulation patterns. Metabolites derived from alpha-linolenic acid are n-3 polyunsaturated fatty acids (PUFAs) (where the double bond is three atoms away from the terminal methyl

group) and metabolites derived from linoleic acid are n-6 PUFAs (double bond is six atoms away from the methyl group). Linoleic acid feeds into the production of arachidonic acid (C20:4) from which a range of pro-inflammatory eicosanoids are derived, for example, prostaglandins and leukotrienes. n-6 PUFA, such as eicosapentaenoic and docosahexanoic acids (as detected in MAP infections) will inhibit arachidonic metabolism. However, it is very much an over-simplification to suggest that n-3 PUFAs minimise inflammation and n-6 PUFAs promote inflammation. For example, prostaglandin E<sub>2</sub> has both a classic pro-inflammatory and an anti-inflammatory role [31]. This stated, the decreases in docosahexanoic acid (Figure 2, clade 4) at 10 to 12 months post MAP inoculation is followed by increases in leukotriene B<sub>4</sub> and prostaglandins. Thus, could indicate that an initial phase in responses to MAP where anti-inflammatory responses are dominant, but then at later phases, inflammation dominates. The potential roles of n-3/n-6 PUFAs ratio in MAP infections has not been assessed but there may be relevant parallels from *M. tuberculosis* infections. *M. tuberculosis* colonisation was increased in the spleens of guinea pigs fed an n-3 PUFA based diet, compared to n-6 PUFA based or control diets [32, 33]. Similarly, *M. tuberculosis* infected guinea pigs fed an n-3 PUFA based diet exhibited increased *M. tuberculosis* pulmonary colonisation compared to controls [34]. Such studies suggest that the observed increased n-3 PUFAs (cis-8, 11,14,17-eicosatetraenoic acid, eicosapentaenoic acid and docosahexanoic acid) encourage MAP colonisation, possibly linked to an anti-inflammatory role.

Arachidonic acid discourages *M. tuberculosis* growth by stimulating the secretion of TNF- $\alpha$  but eicosapentaenoic acid supported colonisation by inhibiting TNF- $\alpha$  secretion [35]. Anes et al. [36] observed similar effects with *M. avium* whereby n-3 PUFAs stimulated growth. In further support of this, incubating macrophage-like J77A4.1 cells with conjugated docosahexanoic acid suppressed the production of pro-inflammatory cytokines (TNF- $\alpha$ , IL-6 and MCP-1) and co-stimulatory molecules (CD40 and CD86) [37]. Consequently, our findings indicate that levels of early n-3 PUFAs and mycobacterial colonisation increase simultaneously. However, our hypothesis may be too simplistic as *M. tuberculosis* infected mice on an n-3 PUFA enriched diet demonstrated elevated pathogen killing [35] and those with a deficient diet showed reduced levels of pro-inflammatory cytokines (IL-6, IL-1 $\alpha$ , MCP1, MIP1 and RANTES) and elevated levels of anti-inflammatory cytokines (IL-4 and growth factor GM-CSF) [38]. Clearly, further MAP-focused studies are required to define the role of the early n-3 PUFA increases. Previous studies suggested that lipid-related metabolomic [14] and lipidomic [39] changes were

indicative of lipid mobilisation that occurred in response to a reduced absorptive capability of the small intestine. However, this is an unlikely cause of our observed lipid-related changes as the heifer calves were asymptomatic throughout the study.

n-6 PUFA increases via the arachidonic acid-derived eicosanoid pathway were one of the most prominent subclasses of metabolites changing on MAP infection (Additional file 2). Many important eicosanoids increased 11 months post MAP inoculation although carbocyclic thromboxane A<sub>2</sub> appear to be increasing at earlier time point (Figure 2). Crucially, two of these metabolites, leukotriene B<sub>4</sub> and bicyclo-PGE<sub>2</sub> were identified in our previous study underlying their validity as biomarkers [16]. Eicosanoids processing from arachidonic acid can follow one of two enzymatic routes; the cyclooxygenase (COX) mediated route produces prostaglandins and thromboxane's, and the lipoxygenase (LOX) mediated route produces hydroxyeicosatetraenoic acids (HETEs) and leukotrienes [40]. Our analyses showed that the metabolite end-points of both routes accumulated at later points of MAP infections (Figure 3). As these are mostly involved in mediating inflammation [41], this was consistent with inflammation dominating MAP infection after 11 months, probably after the initial phase where MAP was suppressed. Steps which could be suppressed could include neutrophil migration and recruitment of CD4 and CD8 T lymphocytes which have been linked to LOX-derived metabolites promotion [42, 43]. There is some evidence for this from transcriptomic assessments where the LOX pathway, as indicated by ALOX15, is downregulated in both MAP positive cows [44] and calves 6 months post MAP inoculation [45]. Further, ALOX5 was down-regulated in primary monocyte-derived macrophages in MAP positive cows [46]. This would align with the delayed accumulation of LOX-dependent metabolites, including 5-HETE, 10, 11-dihydro-leukotriene B<sub>4</sub> and leukotriene B<sub>4</sub> in our metabolomic study (Figure 2).

Considering the COX pathway, it seems likely that this does not wholly conform to the pro-inflammatory phase model in our MAP infection experiment. The literature suggests that expression of the COX2 encoding gene, PTGS2, is rapidly increased in primary monocyte-derived macrophages from MAP positive cows at 8 to 24 h post-infection [46]. Outside of MAP studies, *M. bovis* elicits elevated levels of COX-dependent prostaglandin E<sub>2</sub> [47], as well as increased expression of prostaglandin E synthase [48]. We did observe an increase in thromboxane A<sub>2</sub> in first few months following MAP infection. Thromboxane A<sub>2</sub> is produced by platelets and vaso-constrictive and prothrombotic properties, so is not directly inflammatory. Further,

thromboxane A2 is unstable so that its production must be persistent to allow its detection in our assays [49]. It may be that early COX metabolism feeds into this sub-pathway, rather than more proinflammatory COX products such as prostaglandin E<sub>1</sub> and bicyclo-PGE2 which were observed at later timepoints of MAP infection. However, we also observed later increases in prostaglandin E<sub>2</sub> which increases levels of the regulatory cytokine IL-10 although decreasing T-cell proliferation and Th1 cytokine production but and immunosuppressive programmed death ligand 1 [50]. The absence of clear mechanistic trends to emerge from our eicosanoid assessments could be as a result of our focusing on sera which represented the pooling of metabolites arising from a range of cell and tissue specific effects. Further research assessing immune system parameters, including in specific cell types, are required to define the influence of LOX-dependent and COX-dependent metabolites on inflammatory events and the immune system.

Metabolomic analysis of experimentally MAP-inoculated and control heifer calves using untargeted FIE-HRMS successfully differentiated between experimental groups between 1- and 19- months post MAP inoculation. Broadly, two main types of response were observed, a rapid increase (within 1 month) of saturated fatty acids and some n-3 PUFA that could promote colonisation and suppress inflammation, and a later accumulation of fatty acids including n-6 PUFA and COX products that could indicate a more pro-inflammatory, immune-modulatory phase. Crucially, six fatty acyls were able to differentiate between experimental groups throughout the study. Moreover, several metabolites that were suggested to be biomarkers for MAP in naturally infected heifer calves were also observed in this study where heifer calves were experimentally inoculated. Future work could include examining immune system parameters alongside these targeted metabolites.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13567-022-01087-0>.

**Additional file 1. Metabolites that significantly change following MAP inoculation.** *P* values are indicated. Details of the identified metabolites along with *p*-values of MAP status, time and MAP status\*time.

**Additional file 2. The sub (classes) and AUC values of metabolites differentially expressed in MAP-inoculated heifer calves at 19-months of age.** Details of the identified metabolites including; metabolite ionisation modes, class, subclass, AUC, *P*-value and Log<sub>2</sub> (FC) at 19-months of age.

**Additional file 3. Significantly enriched pathways in MAP-inoculated heifer calves in the combined ionisation mode.** The output of MSEA using ORA demonstrating the significantly enriched pathways in MAP-inoculated heifer calves in the combined ionisation mode.

**Additional file 4. Box and whisker plots of metabolites which display minimal overlapping between groups, MAP-inoculated and control heifer calves, between 1-month and 19-months of age.** (A) 10-Octadecenoic acid; (B) heptadecanoic acid. Blue boxplots = MAP-inoculated heifer calves, green boxplots = control heifer calves.

**Additional file 5. Box and whisker plots of docosahexaenoic acid which displays minimal overlapping between groups, MAP-inoculated and control heifer calves, between 1-month and 19-months of age.** Blue boxplots = MAP-inoculated heifer calves, green boxplots = control heifer calves.

## Acknowledgements

The authors are grateful to KESS2 for funding this work and Helen Phillips for assisting with the FIE-HRMS analysis, as well as Bernardo Villarreal-Ramos and Hans-Martin Vordermeier for their continued support and advice.

## Authors' contributions

DR, LAJM and APK conceptualized the data. APK designed and conducted the MAP-inoculation study. M.B. developed the FIE-HRMS methodology. ENT conducted metabolomic experiments and analysed data. ENT wrote the original draft. GH, DR, LAJM and APK reviewed and edited the manuscript. All authors read and approved the final manuscript.

## Funding

This research was funded by Knowledge Economy Skills Scholarships (KESS), a pan-Wales higher level skills initiative led by Bangor University on behalf of the HE section in Wales, grant number AU30033. The KESS2 project sponsor was ProTEM Services Ltd with contributions from Sona Nanotech. The APC were funded by Wageningen Bioveterinary Research.

## Availability of data and materials

The datasets generated and/or analysed during the current study are available in the Mendely Data repository [51].

## Declarations

### Ethics approval and consent to participate

The animal experiments were approved by the Animal Welfare Body of Utrecht University (permit number 0202.0806) in accordance with the Dutch regulations on animal experimentation.

### Competing interests

The authors declare that they have no competing interests.

### Author details

<sup>1</sup>Institute of Biological, Environmental and Rural Sciences (IBERS), Aberystwyth University, Ceredigion SY23 3DA, UK. <sup>2</sup>Centre of Excellence for Bovine Tuberculosis, Aberystwyth University, Ceredigion SY23 3DA, UK. <sup>3</sup>ProTEM Services Ltd, Horsham RH12 4BD, West Sussex, UK. <sup>4</sup>Wageningen Bioveterinary Research, 8221 RA Lelystad, The Netherlands. <sup>5</sup>Faculty of Veterinary Medicine, Population Health Systems, Utrecht University, 3584 CS Utrecht, The Netherlands.

Received: 8 April 2022 Accepted: 3 July 2022

Published online: 02 September 2022

## References

- Sweeney RW (1996) Transmission of paratuberculosis. *Vet Clin North Am Food Anim* 12:305–312. [https://doi.org/10.1016/s0749-0720\(15\)30408-4](https://doi.org/10.1016/s0749-0720(15)30408-4)
- Whitlock RH, Buergelt C (1996) Preclinical and clinical manifestations of paratuberculosis (including pathology). *Vet Clin North Am Food Anim Pract* 12:345–356. [https://doi.org/10.1016/s0749-0720\(15\)30410-2](https://doi.org/10.1016/s0749-0720(15)30410-2)
- Behr MA, Collins DM (2010) Paratuberculosis: organism, disease, control. CABi, Wallingford
- Rasmussen P, Barkema HW, Mason S, Beaulieu E, Hall DC (2021) Economic losses due to John's disease (paratuberculosis) in dairy cattle. *J Dairy Sci* 104:3123–3143. <https://doi.org/10.3168/jds.2020-19381>

5. Salem M, Heydel C, El-Sayed A, Ahmed SA, Zchock M, Baljer G (2013) *Mycobacterium avium* subspecies *paratuberculosis*: an insidious problem for the ruminant industry. *Trop Anim Health Prod* 45:351–366. <https://doi.org/10.1007/s11250-012-0274-2>
6. Whitlock RH, Wells SJ, Sweeney RW, Van Tiem J (2000) ELISA and faecal culture for paratuberculosis (Johne's disease): sensitivity and specificity of each method. *Vet Microbiol* 77:387–398. [https://doi.org/10.1016/S0378-1135\(00\)00324-2](https://doi.org/10.1016/S0378-1135(00)00324-2)
7. Whittington R, Donat K, Weber F, Nielson S, Eisenberg S, Arrigoni N, Juste R, Sáez JL, Dhand N, Santi A, Michel A, Barkema H, Kralik P, Kostoulas P, Citer L, Griffin F, Barwell R, Moreira A, Slana I, Koehler H, Singh V, Yoo S, Chávez-Gris G, Goodridge A, Ocepek M, Garrido J, Stevenson K, Collins M, Alonso B, Cirone K et al (2019) Control paratuberculosis: who, why and how. A review of 48 countries. *BMC Vet Res*. 15:198. <https://doi.org/10.1186/s12917-019-1943-4>
8. Wells SJ, Collins MT, Faaborg KS, Wees C, Tavornpanich S, Petrini KR, Collins JE, Cernicchiaro N, Whitlock RH (2006) Evaluation of a rapid fecal PCR test for detection of *Mycobacterium avium* subspecies *paratuberculosis* in dairy cattle. *Clin Vaccine Immunol* 13:1125–1130. <https://doi.org/10.1128/CVI.00236-06>
9. Huda A, Jungerson G, Lind P (2004) Longitudinal study of interferon-gamma, serum antibody and milk antibody responses in cattle infected with *Mycobacterium avium* subsp. *paratuberculosis*. *Vet Microbiol* 104:43–53. <https://doi.org/10.1016/j.vetmic.2004.08.011>
10. Suravajhala P, Kogelman LJA, Kadarmideen HA (2016) Multi-omic data integration and analysis using systems genomics approaches: methods and applications in animal production, health and welfare. *Genet Sel Evol* 43:38. <https://doi.org/10.1186/s12711-016-0217-x>
11. Bay S, Begg D, Ganneau C, Branger M, Cochard T, Bannantine JP, Kohler H, Moyer JL, Whittington RJ, Biet FB (2021) Engineering synthetic lipopeptide antigen for specific detection of *Mycobacterium avium* subspecies *paratuberculosis* infection. *Front Vet Sci* 8:637841. <https://doi.org/10.3389/fvets.2021.637841>
12. Karuppusamy S, Mutharia L, Kelton D, Plattner B, Mallikarjunappa S, Karrow N, Kirby G (2021) Detection of *Mycobacterium avium* subspecies *paratuberculosis* (MAP) microorganisms using antigenic MAP cell envelope proteins. *Front Vet Sci* 8:615029. <https://doi.org/10.3389/fvets.2021.615029>
13. Kiser JN, Wang Z, Zanella R, Scraggs E, Neupane M, Cantrell B, van Tassel CP, White SN, Taylor JF, Neibergs HL (2021) Functional variants surrounding endotoxin 2 are associated with *Mycobacterium avium* subspecies *paratuberculosis* infection. *Front Vet Sci* 8:625323. <https://doi.org/10.3389/fvets.2021.625323>
14. de Buck J, Shaykhutdinov R, Barkema HW, Vogel HJ (2014) Metabolomic profiling in cattle experimentally infected with *Mycobacterium avium* subsp. *paratuberculosis*. *PLoS One* 9:e111872. <https://doi.org/10.1371/journal.pone.0111872>
15. Tata A, Pallante I, Massaro A, Miano B, Bottazzari M, Fiorini P, Dal Pra M, Paganini L, Stefani A, de Buck J, Piro R, Pozzato N (2021) Serum metabolomic profiles of paratuberculosis infected and infectious dairy cattle by ambient mass spectrometry. *Front Vet Sci* 7:625067. <https://doi.org/10.3389/fvets.2020.625067>
16. Taylor EN, Beckmann M, Villareal-Ramos B, Vordermeier HM, Hewinson G, Rooke D, Mur LAJ, Koets AP (2021) Metabolomic changes in naturally MAP-infected Holstein-Friesian heifers indicate immunologically related biochemical reprogramming. *Metabolites* 11:727. <https://doi.org/10.3390/metabo11110727>
17. Koets A, Hoek A, Langelaar M, Overdijk M, Santema W, Franken P, van Eden W, Rutten V (2006) Mycobacterial 70 kD heat-shock protein is an effective subunit vaccine against bovine paratuberculosis. *Vaccine* 24:2550–2559. <https://doi.org/10.1016/j.vaccine.2005.12.019>
18. Beckmann M, Parker D, Enot DP, Duval E, Draper J (2008) High-throughput, nontargeted metabolite fingerprinting using nominal mass flow injection electrospray mass spectrometry. *Nat Protoc* 3:486–504. <https://doi.org/10.1038/nprot.2007.500>
19. Chong J, Wishart D, Xai J (2019) Using MetaboAnalyst 4.0 for comprehensive and integrative metabolomics data analysis. *Curr Protoc Bioinform*. 68:e86. <https://doi.org/10.1002/cpbi.86>
20. DIMEdb. <https://dimedb.ivers.aber.ac.uk/>. Accessed 28 Oct 2021
21. Foroutan A, Fitzsimmons C, Mandal R, Piri-Moghadam H, Zheng J, Guo AC, Li C, Guan LL, Wishart DS (2020) The bovine metabolome. *Metabolites* 10:223. <https://doi.org/10.3390/metabo10060233>
22. Kravitz A, Pelzer K, Sriranganathan N (2021) The paratuberculosis paradigm examined: a review of host genetic resistance and innate immune fitness in *Mycobacterium avium* subsp. *paratuberculosis* infection. *Front Vet Sci* 8:721706. <https://doi.org/10.3389/fvets.2021.721706>
23. Fecteau M-E, Pitta DW, Vecchiarelli B, Indugu N, Kumar S, Gallagher SC, Fyock TL, Sweeney RW (2016) Dysbiosis of the fecal microbiota in cattle infected with *Mycobacterium avium* subsp. *paratuberculosis*. *PLoS One* 11:e0160353. <https://doi.org/10.1371/journal.pone.0160353>
24. Kaevaska M, Videnska P, Sedlar K, Bartejsova I, Kralova A, Slana I (2016) Faecal bacterial composition in dairy cows shedding *Mycobacterium avium* subsp. *paratuberculosis* in faeces in comparison with nonshedding cows. *Can J Microbiol* 62:538–541. <https://doi.org/10.1139/cjm-2015-0814>
25. Derakhshani H, Tun HM, Cardoso FC, Plaizier JC, Khafipour E, Looor JJ (2017) Linking periparturient dynamics of ruminal microbiota to dietary changes and production parameters. *Front Microbiol* 7:2143. <https://doi.org/10.3389/fmicb.2016.02143>
26. Umanets A, Dinkla A, Vastenhouw S, Ravesloot L, Koets AP (2021) Classification and prediction of *Mycobacterium avium* subsp. *paratuberculosis* (MAP) shedding severity in cattle based on young stock heifer faecal microbiota composition using random forest algorithms. *Anim Microbiome* 3:78. <https://doi.org/10.1186/s42523-021-00143-y>
27. Purdie AC, Plain KM, Begg DJ, de Silva K, Whittington RJ (2019) Gene expression profiles during subclinical *Mycobacterium avium* subspecies *paratuberculosis* infection in sheep can predict disease outcome. *Sci Rep* 9:8245. <https://doi.org/10.1038/s41598-019-44670-w>
28. McCarthy C (1974) Effect of palmitic acid utilization on cell division in *Mycobacterium avium*. *Infect Immun* 9:363–372
29. McCarthy C (1971) Utilization of palmitic acid by *Mycobacterium avium*. *Infect Immun* 4:199–204
30. Pang Q, Zhang T, Wang Y, Kong W, Guan Q, Yan X, Chen S (2018) Metabolomics of early stage plant cell-microbe interaction using stable isotope labelling. *Front Plant Sci* 7:790. <https://doi.org/10.3389/fpls.2018.00760>
31. Scher JU, Pillinger MH (2008) The anti-inflammatory effects of prostaglandins. *J Investig Med* 57:6. <https://doi.org/10.2310/JIM.0b0113e31819aaa76>
32. Paul KP, Leichsenring M, Pfisterer M, Mayatepek E, Wagner D, Domann M, Sonntag HG, Bremer HJ (1997) Influence of n-6 and n-3 polyunsaturated fatty acids on the resistance to experimental tuberculosis. *Metabolism* 46:619–624. [https://doi.org/10.1016/S0026-0495\(97\)90003-2](https://doi.org/10.1016/S0026-0495(97)90003-2)
33. Mayatepek E, Paul K, Leichsenring M, Pfisterer M, Wagner D, Domann M, Sonntag HG, Bremer HJ (1994) Influence of dietary (n-3) poly-unsaturated fatty acids on leukotriene B4 and prostaglandin E2 synthesis and course of experimental tuberculosis in guinea pigs. *Infection* 2:46–52. <https://doi.org/10.1007/BF01739016>
34. McFarland CT, Fan YY, Chapkin RS, Weeks BR, McMurray DN (2008) Dietary polyunsaturated fatty acids modulate resistance to *Mycobacterium tuberculosis* in guinea pigs. *J Nutr* 138:2123–2128. <https://doi.org/10.3945/jn.108.093740>
35. Jordao L, Lengeling A, Bordat Y, Boudou F, Ficquel B, Neyrolles O, Becker PD, Guzman CA, Griffiths G, Anes E (2008) Effects of omega-3 and -6 fatty acids on *Mycobacterium tuberculosis* in macrophages and in mice. *Microbes Infect* 10:1379–1386. <https://doi.org/10.1016/j.micinf.2008.08.004>
36. Anes E, Kuhnlel MP, Bos E, Moniz-Pereira J, Habermann A, Griffiths G (2003) Selected lipids active phagosome actin assembly and maturation resulting in killing of pathogenic mycobacteria. *Nat Cell Biol* 5:793–803. <https://doi.org/10.1038/ncb1036>
37. Bonilla DL, Ly LH, Fan YY, Chapkin RS, McMurray DN (2010) Incorporation of a dietary omega-3 fatty acid impairs murine macrophage responses to *Mycobacterium tuberculosis*. *PLoS One* 5:e10878. <https://doi.org/10.1371/journal.pone.0010878>
38. Hayford FEA, Dolman RC, Ozturk M, Nienaber A, Ricci C, Loots DT, Brombacher F, Blauuw R, Smuts CM, Parihar SP, Malan L (2021) Adjunct n-3 long-chain polyunsaturated fatty acid treatment in tuberculosis reduces inflammation and improves anemia of infection more in C3HeB/FeJ mice with low n-3 fatty acid status than sufficient n-3 fatty acid status. *Front Nutr* 8:695452. <https://doi.org/10.3389/fnut.2021.695452>
39. Wood PL, Erol E, Hoffsis GF, Steinman M, DeBuck J (2018) Serum lipiomics of bovine paratuberculosis: disruption of choline-containing



- glycerophospholipids and sphingolipids. *SAGE Open Med.* 6:1–7. <https://doi.org/10.1177/2050312118775302>
40. Brock TG, Peters-Golden M (2007) Activation and regulation of cellular eicosanoid biosynthesis. *Sci World J* 7:1273–1284. <https://doi.org/10.1100/tsw.2007.180>
  41. Funk CD (2001) Prostaglandins and leukotrienes: advances in eicosanoid biology. *Science* 294:1871–1875. <https://doi.org/10.1126/science.294.5548.1871>
  42. Tager AM, Bromley SK, Medoff BD, Islam SA, Bercury SD, Friedrich EB, Carafone AD, Gerszten RE, Luster AD (2003) Leukotriene B4 receptor BLT1 mediates early effector T cell recruitment. *Nat Immunol* 4:982–990. <https://doi.org/10.1038/ni970>
  43. Coffey MJ, Phare SM, Peters-Golden M (2004) Role of leukotrienes in killing of *Mycobacterium bovis* by neutrophils. *Prostaglandins Leukot Essent Fatty Acids* 71:185–190. <https://doi.org/10.1016/j.plefa.2004.03.012>
  44. Malvisi M, Palazzo F, Morandi N, Lazzari B, Williams JL, Pagnacco G, Minozzi G (2016) Responses of bovine innate immunity to *Mycobacterium avium* subspecies *paratuberculosis* infection revealed by changes in gene expression and levels of microRNA. *PLoS ONE* 11:e0164461. <https://doi.org/10.1371/journal.pone.0164461>
  45. David J, Barkema H, Guan LL, De Buck J (2014) Gene-expression profiling of calves 6 and 9 months after inoculation with *Mycobacterium avium* subspecies *paratuberculosis*. *Vet Res* 45:96. <https://doi.org/10.1186/s13567-014-0096-5>
  46. Ariel O, Gendron D, Dudemaine PL, Gevry N, Ibeagha-Awemu EM, Bissonnette N (2020) Transcriptome profiling of bovine macrophages infected by *Mycobacterium avium* subspecies *paratuberculosis* depicts foam cell and innate immune tolerance phenotypes. *Front Immunol* 10:2874. <https://doi.org/10.3389/fimmu.2019.02874>
  47. Goto S, Konnai S, Hirano Y, Kohara J, Okagawa T, Maekawa N, Sajiki Y, Watari K, Minato E, Kobayashi A, Gondaira S, Higuchi H, Koiwa M, Tajima M, Taguchi E, Uemura R, Yamada S, Kaneko MK, Kato Y, Yamamoto K, Toda M, Suzuki Y, Murata S, Ohashi K (2020) Upregulation of PD-L1 expression by prostaglandin E 2 and the enhancement of IFN- $\gamma$  by anti-PD-L1 antibody combined with a COX-2 inhibitor in *Mycoplasma bovis* infection. *Front Immunol* 7:12. <https://doi.org/10.3389/fvets.2020.00012>
  48. Cheng Y, Chou CH, Tsai HJ (2015) In vitro gene expression profile of bovine peripheral blood mononuclear cells in early *Mycobacterium bovis* infection. *Exp Ther Med* 10:2021–2118. <https://doi.org/10.3892/etm.2015.2814>
  49. Hamberg M, Svensson J, Samuelsson B (1975) Thromboxanes: a new group of biologically active compounds derived from prostaglandin endoperoxides. *Proc Natl Acad Sci U S A* 72:2994–2998. <https://doi.org/10.1073/pnas.72.8.2994>
  50. Sajiki Y, Konnai S, Okagawa T, Nishimori A, Maekawa N, Goto S, Ikebuchi R, Nagata R, Kawaji S, Kagawa Y, Yamada S, Kato Y, Nakajima C, Suzuki Y, Murata S, Mori Y, Ohashi K (2018) Prostaglandin E2 induction suppresses the Th1 immune responses in cattle with Johne's Disease. *Infect Immun* 86:e00910–e917. <https://doi.org/10.1128/IAI.00910-17>
  51. Mendeley Data. <https://data.mendeley.com/datasets/hvh44pvfvp/1>

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

