

REVIEW

Open Access



Multifunctional roles of leader protein of foot-and-mouth disease viruses in suppressing host antiviral responses

Yingqi Liu^{1,2†}, Zixiang Zhu^{1†}, Miaotao Zhang² and Haixue Zheng^{1*}

Abstract

Foot-and-mouth disease virus (FMDV) leader protein (L^{pro}) is a papain-like proteinase, which plays an important role in FMDV pathogenesis. L^{pro} exists as two forms, Lab and Lb, due to translation being initiated from two different start codons separated by 84 nucleotides. L^{pro} self-cleaves from the nascent viral polyprotein precursor as the first mature viral protein. In addition to its role as a viral proteinase, L^{pro} also has the ability to antagonize host antiviral effects. To promote FMDV replication, L^{pro} can suppress host antiviral responses by three different mechanisms: (1) cleavage of eukaryotic translation initiation factor 4 γ (eIF4G) to shut off host protein synthesis; (2) inhibition of host innate immune responses through restriction of interferon- α/β production; and (3) L^{pro} can also act as a deubiquitinase and catalyze deubiquitination of innate immune signaling molecules. In the light of recent functional and biochemical findings regarding L^{pro} , this review introduces the basic properties of L^{pro} and the mechanisms by which it antagonizes host antiviral responses.

Table of Contents

- 1 Introduction
- 2 Different forms of FMDV L^{pro}
- 3 Cleavage activity of L^{pro}
- 4 Cleavage of host proteins induced by L^{pro}
- 5 Suppression of IFN production mediated by L^{pro}
- 6 Deubiquitination activity of L^{pro}
- 7 L^{pro} counteracts innate immune responses through its DUB activity
- 8 A putative SAP domain identified in L^{pro}
- 9 The SAP domain is important for L^{pro} activity
- 10 Conclusions

1 Introduction

Foot-and-mouth disease (FMD) is a highly contagious disease caused by foot-and-mouth disease virus (FMDV). Outbreaks of FMD spread rapidly and usually cause devastating economic losses and trade embargoes. FMDV primarily infects cloven-hoofed animals including cattle, swine, sheep, and various ruminants. The virus belongs to the genus *Aphthovirus* in the *Picornaviridae* family and has seven serotypes: O, A, C, SAT1, SAT2, SAT3, and Asia1. There is poor cross-protection among these serotypes [1].


The genome of FMDV consists of a single-stranded positive-sense RNA with a length of about 8500 nucleotides. The genomic structure can be artificially divided into three parts: the 5' untranslated region (UTR), the open reading frame (ORF), and 3'-UTR. The single long ORF of viral RNA encodes a polyprotein that is subsequently processed into four mature structural proteins (VP1, VP2, VP3, and VP4) which form the capsid, and about 12 non-structural proteins (L^{pro} , 2A, 2B, 2C, 3A, 3B, 3C, 3D, 3AB or 3ABC, 2BC, and 3CD) (Figure 1) [2].

FMDV leader protein (L^{pro}) and 3C pro proteins have proteinase activity [3, 4], and are suggested to have the

*Correspondence: haixuezheng@163.com

[†]Yingqi Liu and Zixiang Zhu contributed equally to this work

¹ State Key Laboratory of Veterinary Etiological Biology, OIE/National Foot and Mouth Diseases Reference Laboratory, Key Laboratory of Animal Virology of Ministry of Agriculture, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou, Gansu, China
Full list of author information is available at the end of the article



ability to inhibit the functions of a variety of host proteins, suppressing cellular immune responses [5–8]. For instance, 3C^{pro} and L^{pro} can induce the cleavage of host eukaryotic translation initiation factor 4γ (eIF4G), limiting the synthesis of various host proteins [7, 9]. This could possibly include type I interferons (IFNs), indirectly promoting viral replication [10]. 3C^{pro} can also cleave the nuclear factor kappa B (NF-κB) essential modulator (NEMO) and karyopherin α1 (KPNA1) to abate innate immune signaling [5, 6]. Moreover, L^{pro} can directly cleave various other host proteins to suppress antiviral responses [11].

L^{pro}, as a viral proteinase, self-cleaves from the nascent viral polyprotein precursor during FMDV infection and plays an important role in viral pathogenesis. L^{pro} has two different forms (termed Lab and Lb) due to the initiation of translation at two functional AUGs that are separated by 84 nucleotides [12]. However, the Lb AUG is more efficiently used than the Lab site despite translation initiating from the Lab site [13, 14]. Hence, Lb is more abundant than Lab. The complete loss of Lab-coding region of FMDV is reported to be lethal for the virus [15], whereas

the viruses with precisely deleted Lb coding regions (leaderless viruses) were viable and could replicate both in cattle and swine. However, these viruses could not induce any pathological changes and their replicative ability was attenuated [16, 17]. Furthermore, the supernatants of primary cell cultures infected with leaderless viruses possess stronger antiviral activity than the supernatants from wild-type FMDV-infected cells [18]. Recent evidence shows that the nature and extent of the residual leader protein sequences of FMDV precisely lacking the Lb-coding sequence determine different growth characteristics in different host-cell systems [19]. Based on these studies, L^{pro} is thought to have multifunctional roles in viral pathogenicity and is considered an important virulence factor of FMDV.

L^{pro} is known to contribute to virus propagation by suppressing host antiviral activity [20]. L^{pro} has an antagonistic effect on host antiviral responses via at least three mechanisms. The most well-characterized mechanism is the cleavage of eIF4G by L^{pro}, which shuts off host cap-dependent mRNA translation, and IFN translation may be included [7, 21]. Additionally, L^{pro} also directly

suppresses production of IFNs (including type I and type III) at the transcriptional level, through disrupting the IFN signaling pathway to inhibit host innate immune responses [8, 22, 23]. Finally, L^{P_{ro}} can significantly inhibit the activation of some signaling transduction molecules involved in antiviral pathways through its deubiquitination enzyme (DUB) activity [22]. In this review, we discuss the current knowledge of these antagonistic mechanisms of L^{P_{ro}} against host antiviral responses.

2 Different forms of FMDV L^{P_{ro}}

FMDV L^{P_{ro}} shows similarities to the members of the cysteine proteinase family in structure and function [24]. It recognizes the junction sites between L^{P_{ro}} and VP4 and then cleaves itself from the polyprotein [4]. This automatic self-processing makes L^{P_{ro}} the first mature viral protein during FMDV infection. The two forms of L^{P_{ro}} (Lab and Lb) generated have been confirmed in vitro and in vivo [4, 25, 26]. Both these forms of L^{P_{ro}} exhibit the same enzymatic properties [27]. Each of them releases itself from the polyprotein via intermolecular or intramolecular self-cleavage [4, 25]. It is deemed that intramolecular self-processing is more efficient than intermolecular self-processing [28]. Nevertheless, the detailed mechanisms for the production of the two forms of L^{P_{ro}} have not been clearly elucidated. The mechanisms for selection of Lab start site (AUG1) or Lb start site (AUG2) for protein synthesis are complex. Through constructing synthetic fusion genes of AUG1 and AUG2, Belsham determined that before initiation of protein synthesis at AUG2, the ribosomes need to scan past AUG1–AUG2. The two initiation sites can both be used efficiently, whereas internal ribosome entry sites (IRESs) contribute to a slight biased utilization of the Lb site [29]. In a translation system mimicking the translation initiation pattern of the FMDV RNA observed during viral infection, the spacer region between two start codons plays a role in start codon recognition and biases the start codon selection towards the second one to initiate protein synthesis. The utilization of the first start codon depends on its sequence context [30]. Another study showed that the selection of AUG2 does not depend on the assembly of 48S complex formation on the 5' side of AUG1 [31]. A recent study based on previous work presented by Belsham [29] revealed a mechanism involving bias-usage of translation initiation sites of L^{P_{ro}}, suggesting that the poor nucleotide context of the Lab-initiation site restricts its translational efficiency. The ribosomes access the Lb site through linear scanning, starting from the upstream IRES proximal to the first initiation codon and this is not an independent entry process [14]. An early study by Poyry et al. suggested an alternative mechanism by which a few ribosomes reach the second initiation site [32].

Mutations in the initiation site of Lb disables the production of progeny viruses in transfected baby hamster kidney (BHK) cells, while mutations in the Lab initiation site do not affect the production of progeny viruses [33]. The precise deletion of the Lb from the A12 strain of FMDV (serotype A) produced viable viruses in BHK cells, while the mutant virus showed a reduced growth rate and produced smaller plaques [15]. A recent report shows that FMDVs (serotype O) lacking complete Lb coding sequences can be obtained in BHK cells by modifying Lab start codons, while the precise deletion of the Lb coding region alone prevents FMDV replication in primary bovine thyroid cells [19]. In addition, the deletion of the “spacer” region between two initiation codons is not lethal for the virus. These findings imply that the L^{P_{ro}} sequence is physiologically associated with FMDV propagation.

Apart from Lab and Lb, another form of L^{P_{ro}} has been observed, which is termed sLb^{P_{ro}} or Lb' [34, 35]. sLb^{P_{ro}} is generated by the removal of six or seven residues from the C-terminal extension (CTE) of L^{P_{ro}} during FMDV infection [36]. The trimming of the CTE of L^{P_{ro}} results in different characteristics of sLb^{P_{ro}}. sLb^{P_{ro}} cannot form homodimers like Lb via interactions of the CTE of one monomer with the substrate-binding site of the neighboring one, and vice versa [34, 35]. The Lb homodimers have been observed by X-ray crystallography and nuclear magnetic resonance (NMR) [34, 35], providing weak evidence for intermolecular reactions during self-cleavage. The X-ray structures of the L protease were obtained with the two forms of the protein, Lb (not Lab) and sLb, which additionally were modified (C51A). However, both the kinetic evidence of cleavage efficiencies and the structural evidence provided by NMR study on the monomeric variant of Lb, have strongly indicated an intramolecular mechanism of self-processing. Moreover, the obvious formation of a homodimer suggests that it may have a potential function in the modulation of enzyme activity; the dimer may be a physiologically active form responsible for the cleavage activities after the self-processing [35, 37]. The loss of the last six or seven residues in the CTE does not affect the cleavage efficiencies of sLb^{P_{ro}} on the eIF4G site. This is because both Lb and sLb^{P_{ro}} use residue C133 and two conserved amino acid residues (D184 and E186) of CTE, mediating binding and cleavage of eIF4GI. However, the cleavage efficiencies of Lb and sLb^{P_{ro}} are different during the intramolecular incision of the polyprotein substrate due to the lack of an intact CTE in sLb^{P_{ro}}, as the presence of at least one intact CTE is more favorable for intermolecular cleavage [38]. Although, the exact role of sLb^{P_{ro}} remains unknown, it is thought to have a function during FMDV infection [38]. A putative SAP domain identified in L^{P_{ro}} is also

involved in the biological activities and functions of L^{Pro}. The mutation in some sites of the SAP domain lead to the production of different forms of L^{Pro}; all with varying functions [39].


3 Cleavage activity of L^{Pro}

L^{Pro}, the first matured protein of FMDV, self-cleaves from the viral genome ORF-encoding polyprotein. The self-release of L^{Pro} is thought to result from both intramolecular [28] and intermolecular [4] cleavage. The sequences of KVQRKLK*GAGQSS at the junction between L^{Pro} and viral structural protein precursor (P1-2A) are thought to be the cleavage sites [4] (Figure 2A). In addition to the self-cleavage activity of L^{Pro}, it can cleave the homologues of host eIF4G in vitro (Figure 2B). The amino acid sequence recognized as the cleavage site of eIF4GI is PSFANLG*RTTLST [40], and VPLLNVG*SRRSQP for eIF4GII [21]. However, there remain some controversies about the precise cleavage sites within eIF4GI and eIF4GII generated by the Lb^{Pro}, because the cleavage sites of eIF4GI or eIF4GII in the virus-infected cells have not been identified.

L^{Pro} is a papain-like cysteine proteinase. Although sequencing shows that L^{Pro} shares low nucleotide identity with papain family members [24], the typically conserved catalytic cysteine and histidine residues belonging to papain-like proteinase have been identified in L^{Pro} [41]. The catalytic cysteine site is located at the top of the central α -helix, and the catalytic histidine site lies opposite to it on a turn between two β -sheets in the right-hand

domain [42]. The most conserved region between papain-like proteases and Lb structures surrounds the active center, particularly the secondary components, α 1 and β 5– β 6 [42].

The crystal structure of L^{Pro} (indicating the Lb^{Pro}) includes a globular domain similar to other members of the papain superfamily cysteine proteinase, and a flexible CTE. L^{Pro} also possesses the same overall folding, which resembles the cellular prototype of papain. However, the pro-peptide binding loop and many other loops found in papain are not observed in L^{Pro} [42]. Members of the papain proteinase superfamily have a corresponding activity unit, which comprises the catalytic triad of Cys/His/Asn [43]. This catalytic unit of Cys/His/Asp is also present in L^{Pro}. According to a detailed comparison of the two active sites, certain hydrogen bonds and water molecules localized at the catalytic site are remarkably conserved. Hydrogen bonds stabilize the side-chain amide group contributing to the oxyanion hole in both enzymes. One of the carboxylate oxygen atoms of Asp164 and amide nitrogen atoms of Asn46 form a hydrogen bond in Lb. In papain, the hydrogen bond comprises a P-Ser176 hydroxyl group and P-Gln19 amide oxygen atom. The multiple discrepancies between the structures of L^{Pro} and cysteine protease give rise to physicochemical differences between the two enzymes. For example, in the soluble state, when the concentration of cations increases, cysteine protease displays excellent tolerance and keeps its original state, whereas the activity of L^{Pro} changes markedly. The fluctuation of pH can significantly



influence the activity of L^{Pro} because its cleavage activity varies greatly in different pH ranges [34].

4 Cleavage of host proteins induced by L^{Pro}

Eukaryotic cellular translation initiation factor 4F (eIF4F) is a protein complex that recruits ribosomes to bind to host mRNA, initiating cap-dependent translation. This recruitment process is a rate-limiting step and therefore regulates translation [44]. The eIF4F complex comprises eIF4E small cap-binding protein, eIF4G scaffolding protein, and eIF4A ATP-dependent RNA helicase with capped-mRNA. The cap binding factor eIF4E, can bind to a segment of eIF4G to facilitate the formation of the eIF4E/cap-mRNA complex. As a core apparatus of eIF4F complex, eIF4G is a scaffolding protein that provides the binding regions for eIF4E, eIF4A, and RNA elements to form the eIF4F complex. The eIF4G protein also provides binding sites that recruit the small ribosomal subunit interacting protein eIF3 (recruiting the 40S ribosomal subunits to the 5'-end of the mRNA in eIF4F complex), poly(A)-binding protein (PABP), and eIF4E kinases Mnk1 (mitogen-activated protein kinase signal-integrating kinase1) and Mnk2, regulating host mRNA translation [45].

eIF4G proteins possess two homologous proteins in yeast, eIF4GI (*TIF4631*) and eIF4GII (*TIF4632*), sharing a similar function. Both of them contain the conserved binding sites for eIF4E, PABP, eIF3 and RNA. For eIF4GI, it is reported that its N-terminal portion provides the binding sites for eIF4E and PABP, whereas eIF4A and eIF3 bind to the C-terminal portion of eIF4GI [46, 47]. Some picornaviruses including poliovirus, human rhinovirus 2, and FMDV can effectively cleave the eIF4GI, yielding N- and C-terminal fragments [40, 47]. FMDV Lb protease can also cleave eIF4GII, generating a C-terminal fragment [48]. The loss of integrity of eIF4GI and eIF4GII blocks the formation of the eIF4F complexes, which directly influences the cellular cap-dependent translation. However, the C-terminal fragment of both eIF4G proteins containing the binding sites for eIF4A and eIF3 can still bind to the FMDV IRES as efficiently as the non-processing eIF4GI and eIF4GII respectively [47, 48]. Studies over the last two decades have shown that regulation of host and viral mRNAs by eIF4G is achieved by different mechanisms. Viral protein synthesis initiated at two distinct sites from artificial fusion genes is independent of the cap-binding eIF4F complex in the presence of IRES [29]. Furthermore, the cleavage products of eIF4GI (C-terminal portion) stimulate the translation of uncapped RNAs and those carrying IRESs [49]. The interaction of the two eIF4G proteins with IRES is an essential event for promoting IRES activity. Therefore, viral RNA translation is unaffected [48, 50].

eIF4GI is a major form of eIF4G, which correlates with inhibition of cellular cap-dependent protein synthesis within FMDV-infected cells [4, 40]. However, cellular protein synthesis can still be maintained at a reduced level, with the complete loss of intact eIF4GI when virus replication is inhibited [51]. The discovery of human eIF4GII, which appears functionally analogous to eIF4GI, has resolved this puzzle [52]. The shut-off of host cell protein synthesis significantly decreases the expression of various cytokines and the major histocompatibility complex (MHC), resulting in delayed host antiviral effects. However, viral uncapped RNA can be translated through an IRES that is independent of intact eIF4G [53]. Therefore, the virus quickly takes over the host machinery to propagate vast numbers of progeny. FMDV lacking L^{Pro} is unable to escape the antiviral response and is not disseminated in the infected animals [16].

Apart from the cleavage of eIF4G, L^{Pro} can cleave a series of cellular proteins, such as eIF3a, polypyrimidine tract-binding protein (PTB), PABP and Gemin5, which are involved in the control of translation, and death domain associated protein (Daxx), a key factor that crosslinks the apoptosis, innate immune responses and transcription control, to interfere with various cellular pathways during viral infection [54]. The events associated with the extent of cytopathic effects in FMDV-infected cells are proteolysis of PTB, which is involved in mRNA stability and RNA localization, interaction of PABP with the entire FMDV 3'-UTR, and the binding of two subunits of eIF3 (eIF3a and b) with the IRES [11]. Recently, Piñeiro et al. [54] reported that the RNA-binding protein Gemin5 is also a target of L^{Pro}. Gemin5 is the RNA-binding factor of a large macromolecule of the survival of motor neuron (SMN) complex, which acts as a down-regulator of cellular mRNA translation and IRES-driven translation initiation [55]. L^{Pro} recognizes the sequence RKAR of Gemin5 and induce its proteolysis, yielding two stable products of molecular weight 85 and 57 kDa within FMDV-infected cells [54]. Daxx has also been identified as a substrate of L^{Pro}, and the RRLR motif is the recognition site. Daxx is a ligand of Fas, acting as a multifunctional adaptor protein in the process of apoptosis, innate immune responses, and in transcriptional regulation [56]. The cleavage recognition site for L^{Pro} in PABP1 has not been identified experimentally. The sequence similarity with other L^{Pro} substrates and the molecular weight of the proteolysis product imply this characteristic [11], and it is deduced that a novel motif containing sequence (R)(R/K)(L/A)(R) is a putative target sequence of L^{Pro}. Hence, neuroguidin, an eIF4E and cytoplasmic polyadenylation element binding protein (CPEB) that plays an important role in neuronal development [57], is hypothesized to be a potential target of L^{Pro},

with the target sequence as AKRRALS [54]. Furthermore, eIF3a and b are essential to the assembly of the translation initiation complex, and are associated with PABP and RNA-binding protein PTB. This is involved in mRNA stability and RNA localization and can be proteolysed by FMDV L^{pro}, whereas PABP can be partial cleaved by L^{pro} [11]. All these studies suggest that L^{pro} can cleave various host proteins and has potential multifunctional roles.

Other than these identified substrates of L^{pro}, various IRES-binding factors that are targets of other picornavirus proteases may contribute to understanding the link between these proteins and L^{pro}. These factors include poly(rC)-binding protein 2, Gemin3 (RNA helicase that is a component of the SMN complex), RIG-I (retinoic acid-inducible gene 1; a cytoplasmic RNA helicase that senses viral infection), MAVS (mitochondrial antiviral-signaling protein), TRIF (Toll/interleukin (IL)-1 receptor domain-containing adaptor inducing IFN- β or innate immune adaptor molecules), and the stress granules protein G3BP [58–62].

5 Suppression of IFN production mediated by L^{pro}

FMDV infection triggers the activation of various pattern recognition receptors (PRRs) and induces a series of antiviral responses; with the transcription factor NF- κ B acting as a sensor in response to the general alteration of the cellular environment. After the PRRs recognize the pathogens, the coordinated activation of various transcription factors including NF- κ B, IFN regulatory factor (IRF)3 and IRF7, are initiated to induce early expression of type I IFNs and activate host antiviral responses [63].

PRR-induced signal transduction can activate NF- κ B to translocate into the nucleus through degradation of NF- κ B inhibitor. Nuclear translocation of NF- κ B is followed by its binding to the promoter sequences of many genes to initiate their transcription. The expression of various cytokine genes such as the proinflammatory factors, chemokines, and adherence factors is greatly enhanced to induce antiviral responses [64, 65]. NF- κ B also promotes secretion of IFN- α/β and their binding to corresponding receptors. This activates the JAK/STAT signaling pathway, which subsequently induces the expression of hundreds of IFN- α/β -stimulated genes (ISGs). ISGs are a class of antiviral genes that directly encode antiviral proteins that suppress virus propagation at different stages of the viral replication cycle [66]. It was recently reported that the enhanced expression of ISGs increases antiviral effects on FMDV [67]. IRFs are transcription factors that are pivotal for inducing activation of IFN- α/β during virus infection; IRF3 and IRF7 are crucial for virus-triggered IFN- α/β secretion [68]. IFN- α/β belong to the family of type I IFNs and serve as the first line of host defenses, displaying critical antiviral activity

[69]. In addition, IFN- λ , a type III IFN, possesses IFN-like activity and is suggested to be a potent antiviral factor that is effective against many viruses [70, 71].

FMDV L^{pro} acts as an antagonist of innate immune responses mainly eliciting the IFN- α/β specific antiviral activity at both protein and mRNA levels. The down-regulation of IFN expression at least in part corresponds to the cleavage of eIF4G by L^{pro}. Both genetically engineered FMDV lacking L^{pro} (A12-LLV2) and wild-type FMDV (A12-IC) were observed to induce the production of IFN- α/β mRNAs in secondary cells from susceptible animals. However, the A12-LLV2 mutant induces greater antiviral activity than the wild type as a consequence of failing to shut off the expression of host cell protein, including IFN- α/β [18]. L^{pro} blocks IFN protein synthesis, as well as synthesis of IFN- β mRNA and at least three ISGs mRNAs [10], including double-stranded RNA-dependent protein kinase (PKR) which plays an important role in inhibition of FMDV replication, 2', 5' oligoadenylate synthetase 1 (OAS1) and myxovirus resistance protein 1 (Mx1). Using microarray technology, a transcriptional profile associated with the antiviral responses against FMDV was systematically analyzed. The results suggested that L^{pro} significantly inhibits NF- κ B-dependent gene expression including expression of IFN- β and ISGs during FMDV infection [72]. Furthermore, it was found that during the acute infection phase, levels of type I IFN in the serum from infected animals significantly increased [73]. These studies indicate that type I IFN production is associated with antiviral effects against FMDV infection and is important in antiviral immune regulation. L^{pro} as a critical virulence factor of FMDV is capable of using multiple strategies to suppress the production of IFNs.

Many picornaviruses have evolutionarily developed subtle strategies that target host factors to subvert IFNs signaling pathways, and survive and replicate in host cells. For example, enterovirus 2A^{pro} counteracts IFNs responses in infected cells by cleaving melanoma differentiation-associated protein 5 (MDA5) and MAVs [74], while the mengovirus utilizes L^{pro} to prevent the production of IFN- α/β by inactivating iron/ferritin-mediated activation of NF- κ B [75]. Cardiovirus L^{pro} induces cellular nuclear transport inhibition by binding to a key trafficking regulator RanGTPase [76].

Accumulating evidence shows that L^{pro} of FMDV inhibits IFN production through interfering with the IFN signaling pathways. De Los Santos et al. determined that L^{pro} can restrict the induction of IFN- β mRNA [10]. The restriction is partially built on the control of transcription factors and their upstream signaling factors by L^{pro}. L^{pro} was shown to be associated with the downregulation of nuclear p65/RelA during FMDV infection [8]. P65/RelA is the core component of NF- κ B, and a decrease

in the integrity of p65/RelA may lead to the reduction of NF- κ B. This ultimately results in downregulation of IFN- β expression and attenuation of host innate immune responses [8]. The mechanism involved in the downregulation of p65/RelA induced by L^{P ρ} remains unclear. Whether the disappearance of p65/RelA is mediated by the cleavage activity of L^{P ρ} has not been confirmed, since no cleavage products of p65/RelA have been determined and no cleavage sites have been mapped until now. Wang et al. observed that L^{P ρ} decreases IRF-3-induced IFN- α/β expression by reducing IRF-3 and IRF-7 expression [77]. L^{P ρ} can also suppress the secretion of IFN- λ 1 by disrupting the IRFs and NF- κ B activation, which is crucial for IFN- λ 1 expression [23]. The strategy adopted by L^{P ρ} is to cut off the connection between the IFN promoters and transcription factors by decreasing the number of transcription factors, thereby inactivating IFN transcription. L^{P ρ} can also use its deubiquitination activity to prevent IFN- α/β production by reducing ubiquitination of several type I IFN signaling molecules (details in next section). All these results indicate that L^{P ρ} uses various strategies to suppress IFN- α/β production and promote FMDV replication.

6 Deubiquitination activity of L^{P ρ}

It is well known that the activation of many signaling events that connect the sensors with the transcription factors are regulated by ubiquitination enzymes. The conjunction of ubiquitin with the signaling molecules contributes to the activation of several of these signaling events [66, 78]. However, there are also deubiquitinating enzymes (DUBs) [79] that can inactivate this complex by cleaving ubiquitin from its substrate proteins [80]. DUBs belong to the proteinase superfamily, of which 100 members have been identified in humans. DUBs can be classified into two main categories, metalloproteases and cysteine proteases [79]. The DUBs such as, A20, cylindromatosis (CYLD) protein, and deubiquitinating enzyme A (DUBA) negatively regulate the ubiquitination process, and hence, are key regulators in antiviral responses. For example, A20 is involved in downregulation of NF- κ B activation, negatively regulating host antiviral responses. A20 is a DUB that can remove K63-linked ubiquitin from the ubiquitinated receptor-interacting protein (RIP) [80]. RIP is a serine/threonine kinase that contains a death domain which can interact with the death receptors Fas and tumor necrosis factor (TNF) receptor 1 to mediate activation of NF- κ B [81]. Deubiquitination of RIP directly abates activation of the NF- κ B signaling pathway [80]. Yokota et al. recently reported that measles virus P protein upregulates A20 to repress Toll-like receptors, inhibiting activation of NF- κ B [82].

Bioinformatics analysis suggests that Lb has a potential DUB structure and conserved DUB catalytic residues


(Cys51 and His148). The observed catalytic residues are highly conserved in the Lb of all seven serotypes of FMDV. Structural analysis indicates that Lb possesses a topology similar to DUB ubiquitin-specific 14 and resembles papain-like protease (PL^{P ρ}) of severe acute respiratory syndrome coronavirus (SARS-CoV) [22, 83]. It has been observed that mutation of the SAP box (I83A/L86A) or the catalytically active site (C51A or D163 N/D164 N) of Lb results in the inactivation of DUB activity of L^{P ρ} [22].

7 L^{P ρ} counteracts innate immune responses through its DUB activity

Over the course of long-term evolutionary processes, many viruses have developed sophisticated strategies to antagonize host antiviral responses. Redirecting the cellular ubiquitination system to suppress innate antiviral immune signaling pathways is one of the strategies. For example, rotavirus NSP1 blocks NF- κ B- and IRF- dependent transcription of type I IFN by inducing proteasome-mediated degradation of IRF3/5/7 or inhibiting I κ B- α (inhibitor of NF- κ B) degradation to prevent NF- κ B activation [84, 85]. The accessory proteins, Viral Protein R and Virion Infectivity Factor of HIV can independently hijack the cellular ubiquitination system to decrease IRF-3 expression through proteasomal degradation and promote virus replication [86]. As a result, the production of host antiviral ISGs and proinflammatory factors is reduced and the antiviral innate responses are attenuated. Moreover, many viruses can hijack host ubiquitination systems to facilitate viral evasion, genomic replication, and exocytosis [87].

In addition to hijacking the host ubiquitination system for virus replication, many viruses have also developed the ability to disrupt cellular ubiquitination machinery to terminate or block several signaling transduction pathways responsible for the induction of antiviral responses [88]. So far, the PL^{P ρ} of several coronaviruses such as, porcine epidemic diarrhea virus, SARS-CoV, and Middle East respiratory syndrome (MERS-CoV) have been shown to possess deubiquitination activity that antagonizes IFN production, indicating that PL^{P ρ} is a multifunctional protein [89, 90]. Similarly, FMDV L^{P ρ} is a papain-like protease that acts as an antagonist of IFN by negatively regulating IFN transcription and IFN mRNA translation [8, 18, 42, 77].

A recent study from Wang et al. has identified a DUB-like activity of Lb of FMDV [22]. It was observed that Lb significantly inhibited ubiquitination of several adaptor signaling molecules of type I IFN pathway, including RIG-I, TBK1, TRAF3, and TRAF6 (Figure 3). The results of sequence alignment and structural bioinformatics analyses indicate that L^{P ρ} and ubiquitin-specific



protease (USP)14 share similar topology [91]. The DUB activity of Lb was further confirmed through observation of the inhibitory effects of Lb on ubiquitination of RIG-I, TRAF3, TRAF6, and TBK1, which eventually prevents activation of the type I IFN pathway. This DUB activity can be abrogated through mutation of the conserved catalytic sites of Lb. The deubiquitinating processes mediated by Lb are similar to those mediated by DUBA and CYLD. Future studies should focus on whether the DUB activity of Lb is involved in the signaling pathways regulated by A20.

8 A putative SAP domain identified in L^{pro}

De Los Santos et al. discovered that FMDV L^{pro} contains a putative SAP domain (scaffold-attachment factor (SAF) A and SAFB, apoptotic chromatin-condensation inducer in the nucleus (ACINUS), and protein inhibitor of activated STAT (PIAS) domain) [39]. SAP is a conserved domain which usually exists in the eukaryotic proteins and involved in nucleic acid binding, DNA metabolism, DNA repair, chromosomal organization, apoptosis, transcriptional regulation, and immune regulation [92].

SMART software analysis of FMDV L^{pro} predicted an SAP domain between amino acids 47 and 83 of Lb. This putative SAP domain in L^{pro} shows >80% amino acid homology with other SAP domains of eukaryotic proteins. Three-dimensional analysis indicates that L^{pro} and the eukaryotic cellular SAP domains share almost the same α -helix-turn- α -helix structure, in which only two amino acid insertions found in the two α -helices of L^{pro} differed from other cellular SAP domains [39]. Furthermore, a motif of IQKL sequence in L^{pro} resembles the LXXLL signature motif that is mostly found in the SAP domain of PIAS. All these observations demonstrate the presence of a putative SAP domain in L^{pro}.

The eukaryotic SAP domain is usually implicated in PIAS-associated functions. The SAP motif in PIAS has been conserved in evolution, from yeast to humans, and this functional motif can recognize and bind to the AT-rich sequence of scaffold/matrix attachment regions (S/MARs) of eukaryotic chromosomes. S/MAR is usually located close to the enhancer sequence so that it provides a special microenvironment for transcription [93]. PIAS is a negative regulator in host antiviral immunity.

For instance, *pias* gene knockout mice show more resistance to bacterial infection and improved antiviral responses to vesicular stomatitis virus. It is proposed that PIAS affects the expression of >60 genes, most of which are cytokine-induced and pathogen-activated genes involved in NF- κ B and STAT signaling pathways. PIAS1 and PIASy are key proteins of the PIAS family and act as inhibitors to negatively regulate NF- κ B- and STAT-dependent gene expression [94]. Furthermore, PIASy adopts distinctive mechanisms to inhibit virus-induced and IFN-stimulated transcription [95]. Intriguingly, some viral proteins are localized in the S/MAR regions, suggesting an interaction between viral proteins and that S/MAR may block host antiviral activities [96]. In addition, there is evidence showing that the VP35 protein of Ebola virus utilizes PIAS to promote sumoylation of IRF7, thus contributing to inhibition of IFN production in immune cells [97]. Until now, whether L^{pro} can adopt an analogous way of using PIAS in inhibiting cellular antiviral activities remains unclear. However, the N-terminal portion of PIAS3 containing the SAP domain was verified to block the NF- κ B activation through binding to the p65/RelA subunit of NF- κ B [98], whether L^{pro} can use this manner to interrupt activation of NF- κ B remains unclear.

9 The SAP domain is important for L^{pro} activity

Zhu et al. found that expression of various IFN-inducible genes, chemokines or transcription factors, especially NF- κ B-dependent gene expression in L^{pro} SAP domain mutant FMDV-infected bovine cells was significantly enhanced compared with the wild-type FMDV-infected cells [72]. De los Santos and his co-workers revealed that SAP domain is a determinant for L^{pro} nuclear subcellular localization. In FMDV-infected cells, L^{pro} progressively translocates to the nucleus, whereas mutation of two residues at positions 55 and 58 of L^{pro} (SAP mutant) significantly prevents nuclear translocation of L^{pro} without affecting the cleavage of eIF4G. This suggests that the SAP domain affects retention of L^{pro} in the nucleus within the FMDV-infected cells. The proper subcellular localization of L^{pro} in the nucleus is deemed to mediate the L^{pro}-dependent degradation of p65/RelA. Observations concerning SAP-related cellular antiviral responses suggest that in SAP-mutant FMDV-infected cells, the mRNA expression levels of several NF- κ B-dependent cytokines, chemokines, and ISGs are higher than in wild-type FMDV-infected cells [39]. Collectively, the aforementioned results demonstrate that subcellular localization of L^{pro} in the nucleus is an important factor in the suppression of innate immune responses, and that the SAP

domain is involved in this process. Besides, a recent study demonstrated that the catalytic activity and SAP domain of L^{pro} were required for suppressing poly(I:C)-induced IFN- λ 1 production [23].

Diaz-San Segundo et al. found that inoculation of pigs with SAP-mutant FMDV (I55A and L58A mutations were introduced in L^{pro}) can induce early protection against FMD [99]. No clinical signs of FMD, viremia, or virus shedding were observed, even when the pigs were inoculated at 100-fold higher doses than those required to cause clinical signs with wild-type FMDV. The SAP-mutant FMDV elicited strong adaptive immune responses that provided complete protection against wild-type FMDV infection. Impressively, the neutralizing antibody response was induced as early as 2 days post-inoculation and lasted for at least 21 days after inoculation. In the blood of pigs inoculated with SAP mutant virus, expression of IFN- α , TNF- α , IL-1, and IL-6 was higher than in pigs inoculated with the wild-type virus. Zhu et al. reported that FMDV manipulates ubiquitin-activating enzyme one to promote viral replication, and the SAP domain of L^{pro} was involved in this process, which indicates that SAP maybe has a novel role [100]. All these studies suggest that FMDV L^{pro} plays an important role in virus replication process, and the SAP domain may be a critical region for the maintenance of the biological activities of L^{pro}.

10 Conclusions

FMDV has evolved numerous strategies to evade host antiviral responses. In order to survive and replicate in host cells, the virus has developed various ways to impair or suppress the induction and activation of antiviral responses, utilizing viral nonstructural proteins. L^{pro} and 3C^{pro} are the main viral factors that antagonize host immune responses, with L^{pro} being one of the most well-characterized proteins. L^{pro} can cleave numerous host proteins, inhibit cellular protein expression, and deubiquitinate some crucial molecules that are essential for the activation of antiviral pathways and signal transduction. Intensive study of FMDV L^{pro} has uncovered several mechanisms by which FMDV replicates in host cells and suppresses host antiviral responses utilizing L^{pro} (Table 1). However, these observations represent only the “tip of the iceberg” and several questions regarding the different forms of L^{pro} and the pathways involved in L^{pro}-mediated antagonistic effects need to be answered. Further studies are necessary to elucidate these unanswered questions and the multifunctional role of L^{pro} in FMDV infection.

Table 1 The target proteins and the multifunctional role of L^{pro}.

Biological functions of L ^{pro}	Auto cleavage	Cleavage activities	Deubiquitination activity	Unknown
Target proteins	Viral polyprotein [4]	eF4G1 [40]; eF4GII [21]	Daxx [54]	RIG-I, TBK1, TRAF3 and TRAF6 [22]; NF-κB [8]; IRF-3/7 [77]
Recognition site/region	KVQRKLIK ²⁰¹ *GAGQSS	eF4Gh:PSFANILG ⁶⁷⁴ *RTTLST; eF4GII:VPLINVG ⁷⁰⁰ *SRRSQP	Daxx:VLARRLR ³⁶⁰ *ENRSLA (RRLR motif)	Unknown
The effects induced by L ^{pro}	Directing the release of L ^{pro} from the nascent polyprotein	Shutting off cellular translation	Regulation of apoptosis and innate immune antiviral responses	Restricting antiviral activities of IFN-α/β and IFN-λ1
		Control of cellular translation	Impairing innate immune signaling pathways	

Abbreviations

FMD: foot-and-mouth disease; FMDV: foot-and-mouth disease virus; L^{pro}: leader protein; 5'-UTR: 5' untranslated region; 3'-UTR: 3' untranslated region; ORF: the open reading frame; eIF4G: eukaryotic translation initiation factor 4 gamma; NF- κ B: nuclear factor kappa B; NEMO: NF- κ B essential modulator; KPNA1: karyopherin α 1; IFN: interferon; DUB: deubiquitination; NMR: nuclear magnetic resonance; eIF4F: eukaryotic cellular translation initiation factor 4F; PABP: poly(A)-binding protein; Mnk: mitogen-activated protein kinase signal-integrating kinase1; IRES: internal ribosome entry site; MHC: major histocompatibility complex; PTB: polypyrimidine tract-binding protein; Daxx: death-domain associated protein; CPEB: cytoplasmic polyadenylation element binding protein; PCBP: poly(rC)-binding protein; RIG-I: retinoic acid-inducible gene 1; MAVS: mitochondrial antiviral-signaling protein; TRIF: Toll/interleukin (IL)-1 receptor domain-containing adaptor inducing interferon- β or innate immune adaptor molecules; PRRs: pattern recognition receptors; IRF: IFN regulatory factor 3; OAS1: 2', 5' oligoadenylate synthetase 1; RIP: receptor interacting protein; TRAF: TNF receptor-associated factor; TBK1: TANK binding kinase 1; PL^{pro}: papain-like protease; PEDV: porcine epidemic diarrhea virus; SARS-CoV: severe acute respiratory syndrome coronavirus; MERS-CoV: middle East respiratory syndrome; SAP domain: scaffold-attachment factor (SAFA) and SAFB, apoptotic chromatin-condensation inducer in the nucleus (ACINUS), and protein inhibitor of activated STAT (PIAS) domain; S/MARs: AT-rich sequence of scaffold/matrix attachment regions; TNF: tumor necrosis factor.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

YL and ZZ compiled the information, and wrote the manuscript; MZ reviewed and revised the manuscript; HZ conceived the idea and critically reviewed and revised the manuscript. All authors read and approved the final manuscript.

Authors' information

Dr Haixue Zheng is a professor in Lanzhou veterinary research institute, CAAS, China. He has focused on foot-and-mouth disease virus (FMDV) pathogenesis and reverse genetic system studies after doctor graduation. He is studying the FMDV-induced innate immune responses and the viral antagonistic strategies from FMDV.

Acknowledgements

This work was supported by grants from the National Natural Sciences Foundation of China (No. 31302118, 31502042 and 31402179), the Gansu Science Foundation for Distinguished Young Scholars (No. 145RJDA328), the International Atomic Energy Agency (16025/R0), the Project Supported by National Science and Technology Ministry (2015BD12B04) and the Key technologies R&D program of Gansu Province (1302NKDA027).

Author details

¹ State Key Laboratory of Veterinary Etiological Biology, OIE/National Foot and Mouth Diseases Reference Laboratory, Key Laboratory of Animal Virology of Ministry of Agriculture, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou, Gansu, China. ² College of Veterinary Medicine, Northwest A&F University, Yangling, Shaanxi, China.

Received: 24 June 2015 Accepted: 7 October 2015

Published online: 28 October 2015

References

- Ding YZ, Chen HT, Zhang J, Zhou JH, Ma LN, Zhang L, Gu Y, Liu YS (2013) An overview of control strategy and diagnostic technology for foot-and-mouth disease in China. *Viol J* 10:78
- Grubman MJ, Baxt B (2004) Foot-and-mouth disease. *Clin Microbiol Rev* 17:465–493
- Conda-Sheridan M, Lee SS, Preslar AT, Stupp SI (2014) Esterase-activated release of naproxen from supramolecular nanofibres. *Chem Commun (Camb)* 50:13757–13760
- Strebel K, Beck E (1986) A second protease of foot-and-mouth disease virus. *J Virol* 58:893–899
- Du Y, Bi J, Liu J, Liu X, Wu X, Jiang P, Yoo D, Zhang Y, Wu J, Wan R, Zhao X, Guo L, Sun W, Cong X, Chen L, Wang J (2014) 3Cpro of foot-and-mouth disease virus antagonizes the interferon signaling pathway by blocking STAT1/STAT2 nuclear translocation. *J Virol* 88:4908–4920
- Wang D, Fang L, Li K, Zhong H, Fan J, Ouyang C, Zhang H, Duan E, Luo R, Zhang Z, Liu X, Chen H, Xiao S (2012) Foot-and-mouth disease virus 3C protease cleaves NEMO to impair innate immune signaling. *J Virol* 86:9311–9322
- Devaney MA, Vakharia VN, Lloyd RE, Ehrenfeld E, Grubman MJ (1988) Leader protein of foot-and-mouth disease virus is required for cleavage of the p220 component of the cap-binding protein complex. *J Virol* 62:4407–4409
- de Los Santos T, Diaz-San Segundo F, Grubman MJ (2007) Degradation of nuclear factor kappa B during foot-and-mouth disease virus infection. *J Virol* 81:12803–12815
- Belsham GJ, McInerney GM, Ross-Smith N (2000) Foot-and-mouth disease virus 3C protease induces cleavage of translation initiation factors eIF4A and eIF4G within infected cells. *J Virol* 74:272–280
- de Los Santos T, de Avila Botton S, Weiblen R, Grubman MJ (2006) The leader proteinase of foot-and-mouth disease virus inhibits the induction of beta interferon mRNA and blocks the host innate immune response. *J Virol* 80:1906–1914
- Rodriguez Pulido M, Serrano P, Saiz M, Martinez-Salas E (2007) Foot-and-mouth disease virus infection induces proteolytic cleavage of PTB, eIF3a, b, and PABP RNA-binding proteins. *Virology* 364:466–474
- Esteban-Torres M, Landete JM, Reveron I, Santamaria L, de las Rivas B, Munoz R (2015) A *Lactobacillus plantarum* esterase active on a broad range of phenolic esters. *Appl Environ Microbiol* 81:3235–3242
- Gu X, Kumar S, Kim E, Kim Y (2015) A whole genome screening and RNA interference identify a juvenile hormone esterase-like gene of the diamondback moth, *Plutella xylostella*. *J Insect Physiol* 80:81–87
- Poyry TA, Jackson RJ (2011) Mechanisms governing the selection of translation initiation sites on foot-and-mouth disease virus RNA. *J Virol* 85:10178–10188
- Piccione ME, Rieder E, Mason PW, Grubman MJ (1995) The foot-and-mouth disease virus leader proteinase gene is not required for viral replication. *J Virol* 69:5376–5382
- Brown CC, Piccone ME, Mason PW, McKenna TS, Grubman MJ (1996) Pathogenesis of wild-type and leaderless foot-and-mouth disease virus in cattle. *J Virol* 70:5638–5641
- Chinsangaram J, Mason PW, Grubman MJ (1998) Protection of swine by live and inactivated vaccines prepared from a leader proteinase-deficient serotype A12 foot-and-mouth disease virus. *Vaccine* 16:1516–1522
- Chinsangaram J, Piccone ME, Grubman MJ (1999) Ability of foot-and-mouth disease virus to form plaques in cell culture is associated with suppression of alpha/beta interferon. *J Virol* 73:9891–9898
- Belsham GJ (2013) Influence of the Leader protein coding region of foot-and-mouth disease virus on virus replication. *J Gen Virol* 94:1486–1495
- Steinberger J, Skern T (2014) The leader proteinase of foot-and-mouth disease virus: structure-function relationships in a proteolytic virulence factor. *Biol Chem* 395:1179–1185
- Gradi A, Foeger N, Strong R, Svitkin YV, Sonenberg N, Skern T, Belsham GJ (2004) Cleavage of eukaryotic translation initiation factor 4GII within foot-and-mouth disease virus-infected cells: identification of the L-protease cleavage site in vitro. *J Virol* 78:3271–3278
- Wang D, Fang L, Li P, Sun L, Fan J, Zhang Q, Luo R, Liu X, Li K, Chen H, Chen Z, Xiao S (2011) The leader proteinase of foot-and-mouth disease virus negatively regulates the type I interferon pathway by acting as a viral deubiquitinase. *J Virol* 85:3758–3766
- Wang D, Fang L, Liu L, Zhong H, Chen Q, Luo R, Liu X, Zhang Z, Chen H, Xiao S (2011) Foot-and-mouth disease virus (FMDV) leader proteinase negatively regulates the porcine interferon-lambda1 pathway. *Mol Immunol* 49:407–412
- Gorbalenya AE, Koonin EV, Lai MM (1991) Putative papain-related thiol proteases of positive-strand RNA viruses. Identification of rubi- and aphthovirus proteases and delineation of a novel conserved domain

- associated with proteases of rubi-, alpha- and coronaviruses. *FEBS Lett* 288:201–205
25. Clarke BE, Sangar DV, Burroughs JN, Newton SE, Carroll AR, Rowlands DJ (1985) Two initiation sites for foot-and-mouth disease virus polyprotein in vivo. *J Gen Virol* 66:2615–2626
 26. Piccone ME, Diaz-San Segundo F, Kramer E, Rodriguez LL, de los Santos T (2011) Introduction of tag epitopes in the inter-AUG region of foot and mouth disease virus: effect on the L protein. *Virus Res* 155:91–97
 27. Medina M, Domingo E, Brangwyn JK, Belsham GJ (1993) The two species of the foot-and-mouth disease virus leader protein, expressed individually, exhibit the same activities. *Virology* 194:355–359
 28. Glaser W, Cencic R, Skern T (2001) Foot-and-mouth disease virus leader proteinase: involvement of C-terminal residues in self-processing and cleavage of eIF4G1. *J Biol Chem* 276:35473–35481
 29. Greve J, Bas M, Hoffmann TK, Schuler PJ, Weller P, Kojda G, Strassen U (2015) Effect of C1-Esterase-inhibitor in angiotensin-converting enzyme inhibitor-induced angioedema. *Laryngoscope* 125:E198–E202
 30. Crowther M, Bauer KA, Kaplan AP (2014) The thrombogenicity of C1 esterase inhibitor (human): review of the evidence. *Allergy Asthma Proc* 35:444–453
 31. Scozzafava A, Kalin P, Supuran CT, Gulcin I, Alwasel SH (2015) The impact of hydroquinone on acetylcholine esterase and certain human carbonic anhydrase isoenzymes (hCA I, II, IX, and XII). *J Enzym Inhib Med Chem*. doi:10.3109/14756366.2014.999236
 32. Poyry TA, Hentze MW, Jackson RJ (2001) Construction of regulatable picornavirus IRESes as a test of current models of the mechanism of internal translation initiation. *RNA* 7:647–660
 33. Cao X, Bergmann IE, Fullkrug R, Beck E (1995) Functional analysis of the two alternative translation initiation sites of foot-and-mouth disease virus. *J Virol* 69:560–563
 34. Guarne A, Hampoelz B, Glaser W, Carpena X, Tormo J, Fita I, Skern T (2000) Structural and biochemical features distinguish the foot-and-mouth disease virus leader proteinase from other papain-like enzymes. *J Mol Biol* 302:1227–1240
 35. Cencic R, Mayer C, Juliano MA, Juliano L, Konrat R, Kontaxis G, Skern T (2007) Investigating the substrate specificity and oligomerisation of the leader protease of foot and mouth disease virus using NMR. *J Mol Biol* 373:1071–1087
 36. Sangar DV, Clark RP, Carroll AR, Rowlands DJ, Clarke BE (1988) Modification of the leader protein (Lb) of foot-and-mouth disease virus. *J Gen Virol* 69:2327–2333
 37. Steinberger J, Kontaxis G, Rancan C, Skern T (2013) Comparison of self-processing of foot-and-mouth disease virus leader proteinase and porcine reproductive and respiratory syndrome virus leader proteinase nsp1alpha. *Virology* 443:271–277
 38. Steinberger J, Grishkovskaya I, Cencic R, Juliano L, Juliano MA, Skern T (2014) Foot-and-mouth disease virus leader proteinase: structural insights into the mechanism of intermolecular cleavage. *Virology* 468–470:397–408
 39. de los Santos T, Diaz-San Segundo F, Zhu J, Koster M, Dias CC, Grubman MJ (2009) A conserved domain in the leader proteinase of foot-and-mouth disease virus is required for proper subcellular localization and function. *J Virol* 83:1800–1810
 40. Kirchwegger R, Ziegler E, Lamphear BJ, Waters D, Liebig HD, Sommergruber W, Sobrino F, Hohenadl C, Blaas D, Rhoads RE, Skern T (1994) Foot-and-mouth disease virus leader proteinase: purification of the Lb form and determination of its cleavage site on eIF-4 gamma. *J Virol* 68:5677–5684
 41. Nieter A, Haase-Aschoff P, Kelle S, Linke D, Krings U, Popper L, Berger RG (2015) A chlorogenic acid esterase with a unique substrate specificity from *Ustilago maydis*. *Appl Environ Microbiol* 81:1679–1688
 42. Guarne A, Tormo J, Kirchwegger R, Pfistermueller D, Fita I, Skern T (1998) Structure of the foot-and-mouth disease virus leader protease: a papain-like fold adapted for self-processing and eIF4G recognition. *EMBO J* 17:7469–7479
 43. Brocklehurst K, Philpott MP (2013) Cysteine proteases: mode of action and role in epidermal differentiation. *Cell Tissue Res* 351:237–244
 44. Gingras AC, Raught B, Sonenberg N (1999) eIF4 initiation factors: effectors of mRNA recruitment to ribosomes and regulators of translation. *Annu Rev Biochem* 68:913–963
 45. Sonenberg N, Hinnebusch AG (2009) Regulation of translation initiation in eukaryotes: mechanisms and biological targets. *Cell* 136:731–745
 46. Colvin OC, Kransdorf MJ, Roberts CC, Chivers FS, Lorans R, Beauchamp CP, Schwartz AJ (2015) Leukocyte esterase analysis in the diagnosis of joint infection: can we make a diagnosis using a simple urine dipstick? *Skelet Radiol* 44:673–677
 47. Lamphear BJ, Kirchwegger R, Skern T, Rhoads RE (1995) Mapping of functional domains in eukaryotic protein synthesis initiation factor 4G (eIF4G) with picornaviral proteases. Implications for cap-dependent and cap-independent translational initiation. *J Biol Chem* 270:21975–21983
 48. Kashima TG, Inagaki Y, Grammatopoulos G, Athanasou NA (2015) Use of chloroacetate esterase staining for the histological diagnosis of prosthetic joint infection. *Virchows Arch* 466:595–601
 49. Torrello G, Ribeiro de Souza FZ, Carrilho E, Hanefeld U (2015) Xylella fastidiosa esterase rather than hydroxynitrile lyase. *Chembiochem* 16:625–630
 50. Pham H, Santucci S, Yang WH (2014) Successful use of daily intravenous infusion of C1 esterase inhibitor concentrate in the treatment of a hereditary angioedema patient with ascites, hypovolemic shock, sepsis, renal and respiratory failure. *Allergy Asthma Clin Immunol* 10:62
 51. Yeom HJ, Jung CS, Kang J, Kim J, Lee JH, Kim DS, Kim HS, Park PS, Kang KS, Park IK (2015) Insecticidal and acetylcholine esterase inhibition activity of Asteraceae plant essential oils and their constituents against adults of the German cockroach (*Blattella germanica*). *J Agric Food Chem* 63:2241–2248
 52. Sabharwal G, Craig T (2015) Recombinant human C1 esterase inhibitor for the treatment of hereditary angioedema due to C1 inhibitor deficiency (C1-INH-HAE). *Expert Rev Clin Immunol* 11:319–327
 53. Belsham GJ (2009) Divergent picornavirus IRES elements. *Virus Res* 139:183–192
 54. Pineiro D, Ramajo J, Bradrick SS, Martinez-Salas E (2012) Gemin5 proteolysis reveals a novel motif to identify L protease targets. *Nucleic Acids Res* 40:4942–4953
 55. Pacheco A, Lopez de Quinto S, Ramajo J, Fernandez N, Martinez-Salas E (2009) A novel role for Gemin5 in mRNA translation. *Nucleic Acids Res* 37:582–590
 56. Michaelson JS, Leder P (2003) RNAi reveals anti-apoptotic and transcriptionally repressive activities of DAXX. *J Cell Sci* 116:345–352
 57. Jung MY, Lorenz L, Richter JD (2006) Translational control by neuroguadin, a eukaryotic initiation factor 4E and CPEB binding protein. *Mol Cell Biol* 26:4277–4287
 58. Perera R, Daijogo S, Walter BL, Nguyen JH, Semler BL (2007) Cellular protein modification by poliovirus: the two faces of poly(rC)-binding protein. *J Virol* 81:8919–8932
 59. Mukherjee A, Morosky SA, Delorme-Axford E, Dybdahl-Sissoko N, Oberste MS, Wang T, Coyne CB (2011) The coxsackievirus B 3C protease cleaves MAVS and TRIF to attenuate host type I interferon and apoptotic signaling. *PLoS Pathog* 7:e1001311
 60. Almstead LL, Sarnow P (2007) Inhibition of U snRNP assembly by a virus-encoded proteinase. *Genes Dev* 21:1086–1097
 61. Barral PM, Sarkar D, Fisher PB, Racaniello VR (2009) RIG-I is cleaved during picornavirus infection. *Virology* 391:171–176
 62. Dougherty JD, White JP, Lloyd RE (2011) Poliovirus-mediated disruption of cytoplasmic processing bodies. *J Virol* 85:64–75
 63. Honda K, Yanai H, Takaoka A, Taniguchi T (2005) Regulation of the type I IFN induction: a current view. *Int Immunol* 17:1367–1378
 64. Grubman MJ, Moraes MP, Diaz-San Segundo F, Pena L, de los Santos T (2008) Evading the host immune response: how foot-and-mouth disease virus has become an effective pathogen. *FEMS Immunol Med Microbiol* 53:8–17
 65. Stahl MC, Harris CK, Matto S, Bernstein JA (2014) Idiopathic nonhistaminergic angioedema successfully treated with ecallantide, icatibant, and C1 esterase inhibitor replacement. *J Allergy Clin Immunol Pract* 2:818–819
 66. Goodbourn S, Didcock L, Randall RE (2000) Interferons: cell signalling, immune modulation, antiviral response and virus countermeasures. *J Gen Virol* 81:2341–2364
 67. Diaz-San Segundo F, Moraes MP, de Los Santos T, Dias CC, Grubman MJ (2010) Interferon-induced protection against foot-and-mouth disease virus infection correlates with enhanced tissue-specific innate immune cell infiltration and interferon-stimulated gene expression. *J Virol* 84:2063–2077

68. Andersen J, VanScoy S, Cheng TF, Gomez D, Reich NC (2008) IRF-3-dependent and augmented target genes during viral infection. *Genes Immun* 9:168–175
69. Taniguchi T, Takaoka A (2002) The interferon-alpha/beta system in antiviral responses: a multimodal machinery of gene regulation by the IRF family of transcription factors. *Curr Opin Immunol* 14:111–116
70. Ank N, West H, Bartholdy C, Eriksson K, Thomsen AR, Paludan SR (2006) Lambda interferon (IFN-lambda), a type III IFN, is induced by viruses and IFNs and displays potent antiviral activity against select virus infections in vivo. *J Virol* 80:4501–4509
71. Kotenko SV, Gallagher G, Baurin VV, Lewis-Antes A, Shen M, Shah NK, Langer JA, Sheikh F, Dickensheets H, Donnelly RP (2003) IFN-lambdas mediate antiviral protection through a distinct class II cytokine receptor complex. *Nat Immunol* 4:69–77
72. Zhu J, Weiss M, Grubman MJ, de los Santos T (2010) Differential gene expression in bovine cells infected with wild type and leaderless foot-and-mouth disease virus. *Virology* 404:32–40
73. Stenfeldt C, Heegaard PM, Stockmarr A, Tjørnhøj K, Belsham GJ (2011) Analysis of the acute phase responses of serum amyloid A, haptoglobin and type I interferon in cattle experimentally infected with foot-and-mouth disease virus serotype O. *Vet Res* 42:66
74. Feng Q, Langereis MA, Lork M, Nguyen M, Hato SV, Lanke K, Ermdad L, Bhoopathi P, Fisher PB, Lloyd RE, van Kuppeveld FJ (2014) Enterovirus 2Apro targets MDA5 and MAVS in infected cells. *J Virol* 88:3369–3378
75. Zoll J, Melchers WJ, Galama JM, van Kuppeveld FJ (2002) The mengovirus leader protein suppresses alpha/beta interferon production by inhibition of the iron/ferritin-mediated activation of NF-kappa B. *J Virol* 76:9664–9672
76. Bacot-Davis VR, Palmenberg AC (2013) Encephalomyocarditis virus Leader protein hinge domain is responsible for interactions with Ran GTPase. *Virology* 443:177–185
77. Wang D, Fang L, Luo R, Ye R, Fang Y, Xie L, Chen H, Xiao S (2010) Foot-and-mouth disease virus leader proteinase inhibits dsRNA-induced type I interferon transcription by decreasing interferon regulatory factor 3/7 in protein levels. *Biochem Biophys Res Commun* 399:72–78
78. Bibeau-Poirier A, Servant MJ (2008) Roles of ubiquitination in pattern-recognition receptors and type I interferon receptor signaling. *Cytokine* 43:359–367
79. Nijman SM, Luna-Vargas MP, Velds A, Brummelkamp TR, Dirac AM, Sixma TK, Bernards R (2005) A genomic and functional inventory of deubiquitinating enzymes. *Cell* 123:773–786
80. Wertz IE, O'Rourke KM, Zhou H, Eby M, Aravind L, Seshagiri S, Wu P, Wiesmann C, Baker R, Boone DL, Ma A, Koonin EV, Dixit VM (2004) De-ubiquitination and ubiquitin ligase domains of A20 downregulate NF-kappaB signalling. *Nature* 430:694–699
81. Kelliher MA, Grimm S, Ishida Y, Kuo F, Stanger BZ, Leder P (1998) The death domain kinase RIP mediates the TNF-induced NF-kappaB signal. *Immunity* 8:297–303
82. Yokota S, Okabayashi T, Yokosawa N, Fujii N (2008) Measles virus P protein suppresses Toll-like receptor signal through up-regulation of ubiquitin-modifying enzyme A20. *FASEB J* 22:74–83
83. Ratia K, Saikatendu KS, Santarsiero BD, Barretto N, Baker SC, Stevens RC, Mesecar AD (2006) Severe acute respiratory syndrome coronavirus papain-like protease: structure of a viral deubiquitinating enzyme. *Proc Natl Acad Sci U S A* 103:5717–5722
84. Kim S, Kim H, Choi Y, Kim Y (2015) A new strategy for fluorogenic esterase probes displaying low levels of non-specific hydrolysis. *Chemistry* 21:9645–9649
85. Devi L, Pawar RM, Makala H, Goel S (2015) Conserved expression of ubiquitin carboxyl-terminal esterase L1 (UCHL1) in mammalian testes. *Indian J Exp Biol* 53:305–312
86. Okumura A, Alce T, Lubyova B, Ezelle H, Strelbel K, Pitha PM (2008) HIV-1 accessory proteins VPR and Vif modulate antiviral response by targeting IRF-3 for degradation. *Virology* 373:85–97
87. Gale M Jr, Sen GC (2009) Viral evasion of the interferon system. *J Interferon Cytokine Res* 29:475–476
88. Viswanathan K, Fruh K, DeFilippis V (2010) Viral hijacking of the host ubiquitin system to evade interferon responses. *Curr Opin Microbiol* 13:517–523
89. Yang X, Chen X, Bian G, Tu J, Xing Y, Wang Y, Chen Z (2014) Proteolytic processing, deubiquitinase and interferon antagonist activities of Middle East respiratory syndrome coronavirus papain-like protease. *J Gen Virol* 95:614–626
90. Gazzard L, Williams K, Chen H, Axford L, Blackwood E, Burton B, Chapman K, Crackett P, Drobnick J, Ellwood C, Epler J, Flagella M, Gancia E, Gill M, Goodacre S, Halladay J, Hewitt J, Hunt H, Kintz S, Lyssikatos J, Macleod C, Major S, Medard G, Narukulla R, Ramiscal J, Schmidt S, Seward E, Wiesmann C, Wu P, Yee S, Yen I, Malek S (2015) Mitigation of acetylcholine esterase activity in the 1,7-diazacarbazole series of inhibitors of checkpoint kinase 1. *J Med Chem* 58:5053–5074
91. Hu M, Li P, Song L, Jeffrey PD, Chenova TA, Wilkinson KD, Cohen RE, Shi Y (2005) Structure and mechanisms of the proteasome-associated deubiquitinating enzyme USP14. *EMBO J* 24:3747–3756
92. Aravind L, Koonin EV (2000) SAP—a putative DNA-binding motif involved in chromosomal organization. *Trends Biochem Sci* 25:112–114
93. Kipp M, Gohring F, Ostendorp T, van Drunen CM, van Driel R, Przybylski M, Fackelmayer FO (2000) SAF-Box, a conserved protein domain that specifically recognizes scaffold attachment region DNA. *Mol Cell Biol* 20:7480–7489
94. Shuai K (2006) Regulation of cytokine signaling pathways by PIAS proteins. *Cell Res* 16:196–202
95. Kubota T, Matsuoka M, Xu S, Otsuki N, Takeda M, Kato A, Ozato K (2011) PIASy inhibits virus-induced and interferon-stimulated transcription through distinct mechanisms. *J Biol Chem* 286:8165–8175
96. Everett RD, Chelbi-Alix MK (2007) PML and PML nuclear bodies: implications in antiviral defence. *Biochimie* 89:819–830
97. Chang TH, Kubota T, Matsuoka M, Jones S, Bradfute SB, Bray M, Ozato K (2009) Ebola Zaire virus blocks type I interferon production by exploiting the host SUMO modification machinery. *PLoS Pathog* 5:e1000493
98. Jang HD, Yoon K, Shin YJ, Kim J, Lee SY (2004) PIAS3 suppresses NF-kappaB-mediated transcription by interacting with the p65/RelA subunit. *J Biol Chem* 279:24873–24880
99. Diaz-San Segundo F, Weiss M, Perez-Martin E, Dias CC, Grubman MJ, de los Santos T (2012) Inoculation of swine with foot-and-mouth disease SAP-mutant virus induces early protection against disease. *J Virol* 86:1316–1327
100. Zhu Z, Yang F, Zhang K, Cao W, Jin Y, Wang G, Mao R, Li D, Guo J, Liu X, Zheng H (2015) Comparative proteomic analysis of wild-type and SAP domain mutant foot-and-mouth disease virus-infected porcine cells identifies the ubiquitin-activating enzyme UBE1 required for virus replication. *J Proteom Res* 14:4194–4206

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

